Published Monthly. Price $5.00 per Annum.

THE JOURNAL

OF THE

AMERICAN CHEMICAL SOCIETY.

VOL. XX.—NO. 1.

JANUARY, 1898.

EDWARD HART, Editor, J. H. LONG, THOMAS B. OSBORNE,
COMMITTEE ON PAPERS AND PUBLICATIONS.

CONTENTS.

PAGE.

Proceedings ..... (1)-(6)

PAPERS.

A New Form of Discharger for Spark Spectra of Solutions. By L. M. Dennis 1

Vinegar Analysis and Some Characteristics of Pure Cider Vinegar. By Albert W. Smith 3

Root Tubercles in Water Culture. By H. A. Weber 9

Derivatives of Silicon Tetrachloride. By Joseph F. X. Harold 13

Some New Ruthenocyanides and the Double Ferrocyanide of Barium and Potassium. By Jas. Lewis Howe and H. D. Campbell 29

The Caffein Compound in Kola. Part II. Kolatannin. By James W. T. Knox and Albert B. Prescott 34

Note 78

Erratum 79

Books Received 79

REVIEW OF AMERICAN CHEMICAL RESEARCH 1-24

EASTON, PA.:

CHEMICAL PUBLISHING COMPANY.

Entered at the Post-office, Easton, Pa., as Second-class Matter.

Orders for back numbers of the Journal should be addressed to the Librarian, F. E. Dodge, 316 Bowne Ave., Flushing, N. Y.

VOL. XX. [JANUARY, 1898.] NO. 1.

THE JOURNAL

OF THE

AMERICAN CHEMICAL SOCIETY.

A NEW FORM OF DISCHARGER FOR SPARK SPECTRA OF SOLUTIONS.

BY L. M. DENNIS.

Received November 10, 1897.

MANY forms of dischargers for examining the spark spectra of solutions have been described, but all of those with which the writer is acquainted possess disadvantages that seriously impair their value.

To prevent the spattering of the liquid upon the slit of the spectroscope the terminals are, in some cases, surrounded by a glass tube. If a spark of even moderate intensity be used, the liquid which is thrown against the walls of the tube soon cuts off most of the light and obscures the spectrum. It is then necessary to disconnect the discharger and to thoroughly clean the walls of the tube, an operation wasteful both of time and material. Others recommend that the glass tube be abandoned and that free terminals be employed. In this case it becomes necessary to place the spectroscope at some distance from the discharger to avoid the spattering of the liquid againt the slit. Again, if the observations are at all prolonged, the liquid in the discharger is soon used up to such an extent that the lower terminal becomes too dry to give a brilliant spectrum. The discharger must then be refilled.

These difficulties are obviated by the following device which we have been using for the past year. No striking originality

is claimed for it, and the chief excuse for here describing it is furnished by its usefulness.

The lower terminal consists of a platinum wire *x* and the graphite cone *c*. The cone is bored as shown to permit of its being slipped over the end of the platinum wire, and the upper end of the cone is allowed to project slightly above the end *d* of the glass tube *e*. The platinum wire passes through a small glass tube, the upper end of which is fused together about the wire. This tube is held in place by a small cork. The remainder of the apparatus is made of glass tubing of the form as shown. The upper end of *m* is drawn out so that there may be slipped over it a piece of rubber tubing small enough to hold the tube *s*. To fill the apparatus, the tube *s* is removed, the discharger is inclined to the left, and the liquid is run in at the upper end of *m*. *s* is then inserted and pushed down nearly to the bottom of *m*. Upon now bringing the apparatus into the upright position, the liquid in the side arm *e* will rise until on a level with the lower end of *s*. *s* is then carefully drawn up until the liquid in *e* has risen to the point *d*. As the liquid is evaporated by the spark at the terminal *c*, it will be maintained at the level *d* by the entrance of air through the lower end of *s*, and this will continue until the liquid in *m* has fallen until about on a level with *d*.

½ size.

The employment of terminals made from Ceylon graphite was first suggested by Hartley,[1] who used pieces shaped like a wedge and made contact between the platinum wire and the carbon cone by wrapping the wire around the cone. The form here suggested permits of an easy removal of the terminal and its replacement by an electrode of other material. If a platinum terminal is desired, the carbon cone is removed and the glass tube carrying the platinum wire is pushed up until in the proper position.

[1] Trans. Royal Society, 175, 52.

The form of discharger above described leaves the terminals free and exposes the slit to spattering. This difficulty is removed by simply covering the slit with a glass microscope slip of suitable size. We use a slip three inches long by two wide and about one-sixteenth of an inch in thickness, and hold it in place by two rubber bands. If the glass becomes spattered, it can easily be wiped clean or replaced by a clean slip.

The upper terminal, which is not shown in the figure, consists of an insulated platinum wire or a graphite cone, around which the wire is wrapped.

CORNELL UNIVERSITY,
October, 1897.

[CONTRIBUTION FROM THE CHEMICAL LABORATORY OF CASE SCHOOL OF APPLIED SCIENCE. NO. 30.]

VINEGAR ANALYSIS AND SOME CHARACTERISTICS OF PURE CIDER VINEGAR.

BY ALBERT W. SMITH.

Received November 17, 1897.

THE laws of Ohio, copying the food laws of several older states, require, among other things, that apple or cider vinegar to be sold in the state shall contain at least two per cent. of cider vinegar solids. As to the chemical composition of what constitute the solids of pure cider vinegar, very little has been published. The *Analyst* for May, 1891, contains some analyses of pure malt vinegar and the prosecution, in 1893, of several cases in the courts of England, by the food inspectors against dealers in distilled and wood-acid vinegar, attracted considerable attention in that country to the subject of the chemical composition of vinegar and methods of distinguishing from each other the several kinds of vinegar upon the English market. Some valuable data were published by Allen and Moore,[1] especially concerning the character of vinegar made from grain. From the discussion of the question by the Society of Public Analysts[2] it appears that the English vinegar market is largely supplied by products made from distilled alcohol and from malted or unmalted grain. In this country the principal sources of vinegar are from apple cider and from distilled alcohol, the

[1] *Analyst*, 18, 240; and 19, 214.
[2] *Analyst*, 18, 180 and 240; 19, 8 and 33.

latter being usually made from maize and malt. Almost all of this distilled or spirit vinegar, which is originally nearly colorless, is colored by caramel in imitation of the color of cider and malt vinegar.

The chemical distinction between cider and spirit vinegar, when pure, offers little difficulty. The cider vinegar, except that made from early summer apples, is of a dark brown color, contains from two and a half to four per cent. solids and 0.30 to 0.50 per cent. ash, and when old has a pleasant, fragrant odor and characteristic taste. The distilled vinegar contains little solid matter and almost no ash, with little taste or odor besides that imparted by its acetic acid contents. The caramel used in this coloring is usually made by burning or caramelizing glucose or from an aqueous extract of charred grain. Although many manufacturers stoutly contend its use to be to impart flavor and "body" to the vinegar, the palpable and sole object of the coloring process is to enable the unscrupulous dealer to sell the product for "pure cider vinegar" or "pure malt vinegar," which, to my personal knowledge, was the all too common practice in Ohio until the statutes against food adulteration were vigorously and honestly enforced.

The chemical distinction of the source of a sample of vinegar must depend largely upon the quantity and composition of its solid constituents. The organic portion of these solids would seem to offer an excellent opportunity for differentiation, in its maleïc and succinic acids, glycerol and sugar contents, and the accurate determination of these would undoubtedly furnish the most valuable basis for judgment. At present, however, the methods in use for the determination of each of these are so tedious and, in the end, so unsatisfactory, that the mineral contents of the ash afford a more advantageous and practical method for the public analyst.

The ash of cider vinegar differs from that of most other kinds in a number of characteristics, only a few of which have heretofore been pointed out.[1]

The first peculiarity to be noticed is that the ash from cider vinegar commences to volatilize and is fusible at a comparatively

[1] Allen: Commercial Organic Analysis, 1, 389; Cox: *Analyst*, 19, 89.

low temperature. If the ash fuses before all the carbon is burned, particles of this are apt to become enclosed and then to resist further oxidation. To prevent this it is best to evaporate and burn small quantities at a time (not to exceed ten cc.), and to keep the temperature as low as is consistent with complete oxidation. A low red heat must not be exceeded, else loss by volatilization occurs.

A second qualitative difference is the fact that the ash from pure cider furnishes a potassium flame unobscured by sodium light, which is usually not the case with a vinegar containing much added natural water, as imitations usually do.

The ash of cider vinegar differs quantitatively from others so far examined in being quite low in chlorides and sulphates, and high in alkaline carbonates and phosphates. A determination of each of these is quickly and easily made by volumetric methods and the results afford data from which reliable conclusions as to the genuineness of the sample may be drawn. About two-thirds of the phosphates of pure cider vinegar ash are soluble in water. In the ash of other vinegars upon the market a much lower proportion of the total phosphates is soluble in water, and the addition of natural water containing lime or magnesia, to pure cider vinegar, reduces this proportion of soluble phosphates, by converting potassium into alkaline earth phosphates.

The complete analysis may be conveniently carried out as follows: For solids, five to ten grams are evaporated in a flat-bottomed dish and dried to constant weight in a water oven. For total acidity five grams, diluted to about fifty cc., are titrated with standard alkali, using phenolphthalein as indicator. The result is usually reported as acetic acid, without attempting to separate this from the small amounts of other organic acids present.

For ash, ten grams are dried and burned in a crucible, using the before-mentioned precaution of a low temperature. This is weighed and dissolved, and the solution tested qualitatively for color of flame and presence of sulphates and chlorides. Unless these are excessive, as compared with a sample of known purity, they need not be determined quantitatively.

For alkalinity of ash and proportion of phosphates, twenty-five

grams are dried and burned, the ash extracted repeatedly with hot water, and the aqueous solution titrated with standard acid, using methyl orange as indicator. The undissolved residue of ash is then treated with nitric acid, the solution partially neutralized, and the phosphorus in each solution separately precipitated by molybdate solution, the yellow precipitate dissolved in ammonia, reduced with zinc and sulphuric acid, and titrated with standard permanganate solution.

All colored samples should be tested for coloring-matter, which, in imitation vinegars, is most commonly caramel. To test for this, mix five to ten cc. of the sample with about twenty-five cc. paraldehyde, and add alcohol until the three liquids become soluble. After standing twelve to twenty-four hours, the caramel will separate as a sticky, dark-brown precipitate, which, after washing with a little absolute alcohol, has the characteristic bitter taste of caramel, and reduces Fehling's solution freely.

TWENTY-TWO SAMPLES PURE CIDER VINEGAR.

Number.	Acetic acid.	Total solids.	Ash.	No. cc. tenth-normal acid required to neutralize ash from 100 grams vinegar.	Milligrms P_2O_5 in water-solution of ash from 100 gms.	Milligrms P_2O_5 in ash not soluble in water.	Total P_2O_5 in ash from 100 grams vinegar.	Original solids = Solids + 1½ × acetic acid.	Ash per 100 parts original solids.	P_2O_5 milligrms per 100 grams original solids.
	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)
1	5.46	2.27	0.44	43.9	22.7[1]	16.3	39.0[1]	10.46	4.21	373
2	3.29	2.69	0.36	31.2	17.3	8.7	26.0	7.63[2]	4.72	341
3	4.22	3.21	0.36	36.0	18.5	9.2	27.7	9.54	3.77	290
4	3.58	2.91	0.31[2]	28.4[2]	15.6	4.2[2]	19.8[2]	8.28	3.74	239
5	3.74	2.14	0.31	36.0	17.5	7.8	25.3	7.75	4.00	326
6	4.74	2.80	0.35	34.4	18.3	6.8	25.1	9.91	3.53	253
7	4.59	2.86	0.32	32.8	19.5	13.8	33.3	11.74	2.73[2]	284
8	4.19	3.00	0.33	36.8	20.5	8.5	29.0	9.28	3.55	312
9	4.92	4.45[1]	0.38	30.8	17.8	9.5	27.3	11.83	3.21	231
10	4.63	3.30	0.40	47.2	21.3	9.8	31.1	10.24	3.90	303
11	3.24[2]	3.89	0.44	45.2	20.5	10.8	31 3	8.78	5.01	348
12	3.89	3.00	0.33	30.4	22.5	10.5	33.0	8.88	3.72	372
13	4.97	2.40	0.37	44.0	15.5	11.0	26.5	9.85	3.76	269
14	6.55	2.76	0.49	44.8	17.0	4.6	21.6	12.58	3.89	172[2]
15	4.11	2.43	0.36	28.8	21.2	5.6	26.8	8.60	4.19	312
16	7.61[1]	2.97	0.43	55.2[1]	21.0	11.0	32.0	14.38[1]	2.99	223
17	4.00	2.53	0.34	35.2	19.4	11.3	30.7	8.53	3.98	360
18	4.08	2.82	0.33	36.0	13.6[2]	19.4[1]	33.0	8.94	3.69	369
19	4.00	2.75	0.51[1]	45.6	17.5	15.5	33.0	8.75	5.83	377[1]
20	3.80	2.50	0.50	48.8	19.9	9.0	28.0	8.20	6.09[1]	352
21	4.20	2.00[2]	0.39	44.0	19.9	9.0	28.9	8.29	4.70	348
22	4.30	2.50	0.46	40.0	22.5	10.0	22.5	8.95	5.14	365
Aver.	4.46	2.83	0.39	38.8	19.1	10.1	28.6	9.65	4.11	310

[1] Maximum. [2] Minimum.

TWENTY-SEVEN SAMPLES CIDER VINEGAR CONTAINING ADDED WATER OR ADDED SPIRIT VINEGAR SOLD IN OHIO MARKETS.

Number.	Acetic acid.	Total solids.	Ash.	No. cc. tenth-normal acid required to neutralize ash from 100 grams vinegar.	Millig'ms P_2O_5 in water solution of ash from 100 gms.	Millig'ms P_2O_5 in ash not soluble in water.	Total P_2O_5 in ash from 100 gms. vinegar.	Original solids = Solids $+ 14 \times$ acetic acid.	Ash per 100 parts original solids.	P_2O_5 millig'ms per 100 grams original solids.
	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)
1	4.10	2.64	0.53	24.4	none	8.4	8.4	8.79	6.03	95
2	4.21	2.94	0.47	10.8	15.2	8.1	23.3	9.25	5.08	252
3	4.11	2.50	0.28	24.4	4.4	4.0	8.4	8.66	3.23	97
4	3.84	2.35	0.30	9.8	4.4	20.2	24.6	8.11	3.70	303
5	4.31	2.10	0.27	24.0	12.8	8.5	21.3	8.56	3.15	249
6	4.26	2.41	0.27	26.8	15.0	5.0	20.0	8.80	3.07	227
7	4.15	2.35	0.27	29.6	...	...	...	8.58	3.15	...
8	4.07	2.11	0.27	27.6	...	...	...	8.24	3.27	...
9	4.01	2.09	0.26	23.0	12.0	5.0	17.0	8.10	3.21	210
10	3.84	2.06	0.26	26.4	...	...	...	7.82	3.33	...
11	3.43	1.69	0.25	18.8	12.0	12.0	24.0	6.83	3.66	351
12	4.46	2.99	0.25	16.6	6.0	3.0	9.0	9.68	2.58	94
13	3.73	1.21	0.25	1.4	none	7.5	7.5	6.81	3.67	110
14	3.97	1.91	0.24	22.8	5.3	8.8	14.1	6.85	3.50	205
15	4.83	2.20	0.24	21.6	11.4	15.8	27.2	9.44	2.54	288
16	3.68	2.45	0.23	15.6	8.6	6.6	15.2	7.97	2.88	191
17	4.00	1.74	0.22	25.2	5.8	8.4	14.2	7.74	2.84	183
18	3.65	1.42	0.20	17.6	...	...	...	6.89	2.90	...
19	4.70	2.14	0.20	13.4	10.0	10.5	20.5	9.19	2.18	223
20	3.01	3.41	0.20	20.8	...	...	...	7.92	2.53	...
21	3.84	1.90	0.20	27.6	5.2	15.2	20.4	7.66	2.61	266
22	4.10	1.20	0.18	14.4	5.0	3.0	8.0	7.35	2.45	109
23	3.88	1.57	0.18	13.0	...	...	...	7.39	2.43	...
24	3.74	1.41	0.17	9.6	...	...	...	7.02	2.42	...
25	4.62	2.02	0.17	11.2	5.8	3.0	8.0	8.95	1.90	89
26	3.44	1.19	0.14	14.0	...	...	...	6.35	2.20	...
27	4.03	1.64	0.14	7.4	trace	4.5	4.5	7.68	1.83	59
				SAMPLE MADE FROM DRIED APPLES AND GLUCOSE.						
	4.29	3.89	0.25	21.0	...	...	5.5	10.33	2.42	53
				SAMPLE MADE FROM APPLE CIDER AND GRAPE JUICE.						
	4.54	2.77	0.30	34.0	...	...	4.0	9.58	3.13	42
				FOUR SAMPLES MALT VINEGAR.						
Eng.	5.90	2.21	0.28	6.0	...	...	12.5	11.06	2.53	113
Am.	5.16	2.67	0.26	6.0	...	...	9.0	10.41	2.50	86
"	4.40	2.04	0.23	5.6	...	...	9.2	8.64	2.66	106
"	4.01	1.75	0.20	4.4	...	...	9.5	7.76	2.58	122
				SIXTY-FIVE SAMPLES SPIRIT VINEGAR.						
Av..	3.84	0.38	0.06	1.1	none	trace				
Max	5.99	0.78	0.15	5.6	"	"				
Min.	2.87	0.14	0.01	0.0	"	"				

In columns 1, 3, 5, and 8 of the accompanying tables, results are stated as grams per 100 grams of the sample. In column 4 are given the number of cubic centimeters of decinormal acid required to neutralize the filtered water solution of the ash derived from 100 grams of the vinegar. In columns 5 and 6 are the number of milligrams of phosphorus pentoxide (P_2O_5) respectively in the water solution and the water insoluble portion of the ash derived from 100 grams of the sample. Column 7 contains the sum of these two quantities, or the total phosphorus pentoxide. In column 8, what Hehner[1] calls "original solids" are given. These numbers are obtained by multiplying the acetic acid quantities by 1.5 and adding to this the total solids. It is assumed by Hehner that three parts of sugar by alcoholic and acetic fermentation yield two parts of acetic acid, this being approximately the theoretical quantitative relations:

$$\underset{180}{C_6H_{12}O_6} \rightarrow 2C_2H_5OH \rightarrow \underset{120}{2HC_2H_3O_2}.$$

Hence, by increasing the acetic acid in the ratio of two to three, we get approximately the theoretical quantity of sugar from which it was produced. By adding to this the amount of solids still remaining in the vinegar, the quantity of solids present in the unfermented liquid is assumed to be obtained. Of course such an assumption is extremely far from the truth, because, from various losses during fermentation, by volatilization of alcohol and acetic acid, only one-half to three-quarters of the theoretical yield of acid is obtained, and this loss varies greatly in the different processes, and in the same process depends largely upon individual care during each of its steps. Further, during the fermenting and settling periods albuminoid and other constituents settle in considerable quantities, so that such calculations possess little real worth, and, when spirit vinegar is used as an adulterant of cider or malt vinegar, they may become decidedly misleading. The figures in columns 8, 9, and 10 are given principally for the sake of comparison with previous analyses which have been reported in this way. In column 9 the percentages of ash to "original solids," found in column 8, have been given, obtained by dividing 100 times the

[1] *Analyst*, 16, 82.

per cent. of ash as given in column 3 by the per cent. of "original solids." In the same way the milligrams of total phosphorus pentoxide per 100 grams of original solids are reported in column 10.

Of these results, the quantity of ash, the alkalinity of the ash, and amount of soluble phosphates are of considerable uniformity in pure cider vinegar, and these, when considered with the other characteristics above enumerated, serve to differentiate this kind quite sharply from all other commercial varieties and to afford a basis for the approximate estimation of the quantity of spirit vinegar or of water, with which pure cider vinegar may have been adulterated.

[CONTRIBUTIONS FROM THE LABORATORY OF AGRICULTURAL CHEMISTRY, OHIO STATE UNIVERSITY.]

ROOT TUBERCLES IN WATER CULTURE.[1]

BY H. A. WEBER.

Received November 22, 1897

SINCE the discovery by Hellriegel of the connection between root tubercles and the fixation of atmospheric nitrogen by leguminous plants, the literature on this interesting and important question is being constantly enriched by painstaking experiments on the part of scientific investigators all over the world. The subject has thus far been studied only in connection with sterilized soil or sand. It occurred to the writer that it might be possible to produce root tubercles in water culture and thus be able to observe the process better than when the roots of the plants were buried in soil or sand. The work of conducting the experiments about to be described was entrusted to Mr. J. C. Britton, one of the writer's advanced students. The results more than met my expectations, and this preliminary notice is made public for the benefit of those who may feel inclined to employ the method in their investigations.

For the past four or five years the writer has employed in water-culture experiments an apparatus, designed by himself which presents certain advantages over the methods heretofore described. As this apparatus was employed in the experiments

[1] Read before the Columbus Section of the American Chemical Society, Nov. 10, 1897.

under consideration, a brief description of the same may not be out of place.

As will be seen from the accompanying illustration, *a* is an aspirator of four liters capacity; b_1 and b_2 are ordinary salt mouth bottles of two liters capacity, which serve as culture jars; *c* is a small flask with a side tube, as used in fractional distilla-



tion; *d* is a collecting bottle of four liters capacity. The culture jars are fitted with cork disks to hold the plants in the usual manner and also support the thistle tube and the siphons as shown in the illustration.

At the beginning of a vegetation experiment, the flask *c*, the culture jars, and the aspirator, are filled with the nutrient solution. The aspirator is closed with a rubber stopper carrying a glass tube, to the end of which a short piece of rubber tubing is attached. In this manner the solution is conveyed to the thistle tube, which extends to near the bottom of jar b_1, the flow being controlled by means of a Hoffman clamp placed on the rubber tubing. By means of a siphon, the short arm of which extends a short distance below the surface of the solution in jar b_1, and the long arm nearly to the bottom of jar b_2, the solution flows into jar b_2.

Another siphon with one arm extending a short distance

below the surface of the solution in jar b_2, and the other below the side tube in flask c, carries the solution into flask c, from which it runs through the side tube into the collecting bottle. The drop of the solution into the thistle tube is so regulated that the aspirator will empty itself in about twenty-four hours, when the solution in the collecting bottle is transferred back into the aspirator. This is done once a day until the vegetation experiment is completed.

The advantages of this arrangement are :

1. Two seedlings, or if, as the writer has frequently done, another culture jar be inserted between jar b_2 and the flask c, three seedlings can be grown in the same solution at the same time, so that, if an accident should happen to one of the seedlings, the experiment need not necessarily fail.

2. The solution is continually being aerated as it drops into the thistle tube and into the collecting bottle, as well as when it is transferred to the aspirator. By this means the roots of the seedlings are constantly supplied with free oxygen, a condition necessary for healthy growth.

3. Plants can be grown to maturity without removing them to other jars filled with fresh solutions.

4. The solution in the culture jars always remains at the same level, *i. e.*, on a level with the side tube of the flask.

Two solutions were prepared as follows :

1. COMPLETE NUTRIENT SOLUTION (Wolff's Solution).

	Grams.
Bone ash dissolved in nitric acid	20
Potassium nitrate	11
Magnesium sulphate	7
Potassium chloride	3
Ferrous sulphate	Minute quantity

The whole dissolved in distilled water, neutralized with sodium carbonate, and diluted to one liter.

2. COMPLETE NUTRIENT SOLUTION WITHOUT NITROGEN.

	Grams.
Bone ash dissolved in nine cc. sulphuric acid	20.0
Potassium sulphate	9.5
Magnesium sulphate	7.0
Potassium chloride	3.0
Ferrous sulphate	Minute quantity

The whole dissolved in distilled water, neutralized with sodium carbonate, and diluted to one liter. For the growing of plants

the apparatus was filled with a mixture of the respective solutions and distilled water in the proportion of five cc. of the solutions to one liter of water. Trials with fifteen cc., and even ten cc. of the solutions to one liter of water, failed on account of over-nutrition, the seedlings refusing to grow and finally dying.

The plants employed were dwarf peas. The seeds were made to germinate in sterilized sand, moistened with distilled water. When the seedlings were four or five inches in height, they were taken from the sand, the roots washed, and the plants transferred to the culture jars.

Three sets of apparatus, as shown in the illustration, were provided. The first set contained the complete nutrient solution. The second set contained solution No. 2, as described above, without nitrogen.

The third set contained the same solution, but in addition the seedlings before being placed into the jars, were inoculated with tubercle germs, by immersing the roots for a few minutes in a cold water infusion of a soil, in which peas had been grown.

As might be expected, the plants in set No. 1, containing the complete nutrient solution, were healthy and vigorous, and produced flowers and fruit.

The plants in set No. 2 grew rapidly for a time, but were weak and sickly in appearance. In about ten days the leaves began to turn yellow and to show the effect of nitrogen starvation. A few small blossoms finally appeared, but this caused the speedy death of the plants. The roots developed very rapidly, completely filling the jars. They were doubtless in search of nitrogenous food.

The plants in set No. 3 grew normally for about ten days, when they began to show the effect of nitrogen starvation in a marked degree. No tubercles could be observed on the roots. About the thirteenth day the plants began to recuperate, the leaves assumed a normal green color, and from this time on the growth was vigorous and normal. On the fifteenth day the tubercles were first observed, but they had then attained considerable size.

The plants, like those of set No. 1, produced flowers and fruit. Investigations in this direction will be continued.

[CONTRIBUTION FROM THE JOHN HARRISON LABORATORY OF CHEMISTRY, No. 25.]

DERIVATIVES OF SILICON TETRACHLORIDE.[1]

BY JOSEPH F. X. HAROLD.

Received December 1, 1897.

GENERAL REMARKS UPON COMPOUNDS OF THE HIGHER HALOIDS OF THE ELEMENTS OF GROUP IV.

THE review of the reactions of these tetrachlorides furnishes some interesting data from which inferences of importance may be drawn. There are, however, numerous problems still unsolved, and many reactions that yet demand study, before any broad comparison of the behaviors of these elements can be definitely made, or any conclusion as to the influence of the atomic weight or metallic character on the reactivity of their tetrachlorides established. No general rule could be deduced from the data already collected, which would enable us by a logically drawn analogy, to predict their behaviors in certain uninvestigated reactions.

Carbon and silicon, the first two elements of the group, are by their position more or less isolated, and removed from too close a comparison with any of the sub-group elements, and their reactions are best studied with a view to determining or increasing their own already well-developed similarities, without seeking to broaden to too great an extent their relation to other elements of the group. The chlorides of silicon and carbon may be said, however, in a general way, to possess in common with the other group members, a distinctly acid nature, and to exhibit the power of combining with bases to form stable, well-defined compounds. This behavior may be predicted for even the unstudied chlorides of germanium, lead and thorium, since every element thus far investigated has in its tetrachloride form thus deported itself. Ammonia, the substituted ammonia derivatives, such as methylamine, toluidine, urea and amido bodies of a distinctly basic character, may here be said to react similarly with all. With acid amides and bodies of a distinctly acid nature, no such general rule may be adopted, and a series of reactions occur which seem to be conditioned by no rise or fall in atomic weight

[1] Thesis presented to the University of Pennsylvania for the degree of Doctor of Philosophy.

or by position and group relation of the elements. Titanium, one of the elements of the sub-group having a less metallic character, and tin of the opposite sub-group and of a distinctly metallic-nature and a high atomic weight, represent the most reactive of the elements of Group IV, in their tetrachloride form. Other than this, there seems to be little relationship between them, being distinctly separated by such considerations as place them in opposite groups. Titanium, the element of lowest atomic weight of the first sub-group, is in its quadrivalent form most reactive. Its tetrahaloid forms derivatives with hydrocyanic acid, cyanogen chloride, and the oxides of nitrogen. It thus has the power of holding acid as well as basic compounds in combination. Zirconium, on the other hand, the element next in succession to titanium, exhibits no such reactions, and its combinations seem limited to compounds of a basic character, as work now in progress in this laboratory indicates. We might then assume the rule "that in the first sub-group the reactivity decreases with a rise in atomic weight, being highest in titanium," with more or less reason. In the case of the opposite sub-group, however, the paucity of data forbids the formulation of any such law. Tin we know to be exceedingly reactive in its quadrivalent haloids, but germanium, the element which precedes it in the sub-group, whose reactions, if studied, would make some such general statement possible, has been but little investigated with such an end in view.

The subsequent chlorides of thorium and lead are being studied at present, and the determination of their reactions will show the relevance of their weight relationship to their behaviors, and make possible a more comprehensive and thorough comparison of the actions of the tetrachlorides of the fourth periodic group. The accompanying table represents the behavior of the several tetrachlorides toward various reagents. The reactivity is indicated by the letter "*r*," and the failure to form compounds with these reagents by the character "*o*."

	Aniline, etc.	Fatty amines.	Urea, thiourea.	Acid amides.	HCN.	Acetonitrile.	Benzonitrile.	Tolunitrile.	Succinonitrile.	Chlorcyanog'n	S_2Cl_2.	NO_2.	PCl_5.	PCl_3.	NH_3.
CCl_4	r	.	.	.	.	.	.	.	.	.	.	r	.	.	r
$SiCl_4$	r	r	r	0	0	0	0	0	0	0	0	0	0	0	r
$TiCl_4$	r	.	.	.	r	r	r	r	r	r	r	.	r	.	r
$GeCl_4$	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
$ZrCl_4$	r	r	.	.	0	0	0	0	0	0	.	.	.	.	r
$SnCl_4$	r	r	.	.	r	r	r	r	r	0	r	.	r	.	r
$PbCl_4$	r	.	.	.	.	.	.	.	.	.	.	.	.	.	r
$ThCl_4$	r	.	.	.	.	.	.	.	.	.	.	.	.	.	r

THE ACTION OF AROMATIC AMINES AND NITRILES ON SILICON TETRACHLORIDE.

The work was entered upon chiefly with a view to developing some new analogies between the behavior of silicon tetrachloride and that of the tetrachlorides of the other members of Group IV, or any characteristic differences which might prove noteworthy. The hope of the investigation was, that by some well-defined series of reactions observed therein, a new means might be found of isolating the members of the group, in their higher forms of combination, from one another or from members of other groups.

The investigation is all the more demanded, since but little work has been done in this field, the action of nitrils having been studied only with the tetrachlorides of titanium and tin.[1] The action, therefore, of nitrils on silicon tetrachloride was a needed expansion of the subject, and this, together with the actions of the chlorides of sulphur and phosphorus, cyanogen and chloride of cyanogen, nitrogen dioxide and nitroxyl chloride upon silicon tetrachloride, constitute the experimental part of the present research, while the demonstration of the inactivity of the latter, as compared with the behaviors of tin and titanium tetrachlorides, determines its scope.

The second part of this article deals with the reactions between silicon tetrachloride and amines of the benzene series. The behavior of these bodies towards higher haloids of the fourth group has been investigated only in the silicon derivatives, and

[1] Henke: *Ann. Chem.* (Liebig), 106, 281; Shinn: Thesis, University of Pennsylvania, 1895.

here the reaction cannot be said to be determined, since the investigators in this field are at odds in their results, and it is hoped that this work will aid in fixing the exact constitution of the products of the reaction of silicon tetrachloride and aromatic amines.

The silicon tetrachloride was made by the action of dry chlorine on silicon, at a low heat, the resulting vapors being collected in chilled condensers. Any free chlorine was removed from the product by shaking with mercury, and the silicon tetrachloride further purified by fractional distillation, till it exhibited the constant boiling-point, 57°–59°. The silicon used in this preparation was made by the reduction of very fine sand, free from iron, by magnesium powder, according to the method recommended by Gattermann.

THE ACTION OF FORMONITRILE ON SILICON TETRACHLORIDE.

Dry hydrocyanic acid gas conducted into cool silicon tetrachloride produces no change; no product is formed and the silicon tetrachloride boils at the usual temperature. Wöhler[1] had prepared the compound $TiCl_4.2HCN$, while Klein[2] got a corresponding tin derivative, as did also Shinn.[3] These results led to the attempt to combine formonitrile and silicon tetrachloride with the result above indicated.

ACETONITRILE.

Henke[4] had shown that methyl cyanide combined with titanium and tin tetrachlorides, to form derivatives of the type $SnCl_4.2CH_3CN$. Anhydrous acetonitrile, made by the distillation of acetamide and phosphorus pentasulphide, and further purification of the product, was added to silicon tetrachloride chilled in ice; no action was apparent or heat generated. The precaution of chilling in ice was taken, because in the case of tin derivatives the amount of heat produced in the reaction was sufficient to decompose the products. Here, however, even when the ordinary temperature was resumed, no combination was effected. The slightest trace of moisture, however, serves

[1] *Ann. Chem.* (Liebig), **73**, 226.
[2] *Ann. Chem.* (Liebig), **74**, 86.
[3] Thesis, University of Pennsylvania, 1896.
[4] *Ann. Chem.* (Liebig), **106**, 281.

to precipitate a silica compound which retains the nitrile tenaciously. Its analysis, however, indicates no definite constitution, and it is probably nothing more than a mixture of silicic acid and methyl cyanide.

PROPIONITRILE.

This was made by the distillation of potassium cyanide with potassium ethyl sulphate; the isopropionitrile formed at the same time, being removed by a small quantity of concentrated hydrochloric acid. As with acetonitrile, the anhydrous liquid gives no compound with silicon tetrachloride, while the tetrachlorides of tin and titanium afford derivatives of a constitution similar to the acetonitrile bodies. (See above references.)

SUCCINONITRILE.

The nitrile of succinic acid was made by digesting ethylene dibromide with an alcoholic solution of potassium cyanide, and after being rendered absolutely anhydrous was added to silicon tetrachloride. It gave no derivative, being in this distinguished from tin, which forms the compound $SnCl_4.2(CH_2CN)_2$.[1]

AROMATIC NITRILES. BENZONITRILES.

This reagent was made by distilling benzoic acid and dry potassium thiocyanate, adding ammonia to the semi-solid distillate and extracting with ether. With this nitrile the behavior of tin and titanium chlorides had been investigated by Henke,[2] who obtained the compounds $C_6H_5CN.TiCl_4$ and also $C_6H_5CN.SnCl_4$. With silicon tetrachloride, however, it gives no derivative under ordinary conditions or by the action of heat and pressure. Silicon tetrafluoride also exhibits no reaction with this nitrile.

TOLUNITRILE.

The tetrachloride here gives a compound, but neither the chloride nor fluoride of quadrivalent silicon unite with it. No variation in the result was noticed when the silicon tetrachloride was heated under pressure with the nitrile, these compounds remaining uncombined as before.

[1] Shinn: University of Pennsylvania, Thesis, 1896.
[2] *Ann. Chem.* (Liebig), 106, 181.

The *nitriles* of *mandelic* and *lactic acids* gave no derivatives with silicon tetrachloride under ordinary conditions, but when heated under pressure, silicic acid and complex tarry decomposition products resulted, which did not attract analysis. In all the above cases the mediums of interactions were benzene and ether, the latter being preferable, because it was thought the more easily dried.

NITROGEN DIOXIDE.

Marignac[1] obtained the derivative of carbon tetrachloride, to which he assigned the formula $CCl_4.2NO_2$. Later, Kuhlmann[2] prepared derivatives of silicon tetrachloride with every oxide of nitrogen. On passing NO_2, prepared by heating lead nitrate, into silicon tetrachloride, no heat development was noticed. The latter dissolves the gas and becomes colored by the same, but a single distillation, during which the boiling-point showed no change, sufficed to completely decolorize the liquid and remove the nitrogen dioxide from it.

THE MIXED VAPORS OF NO_2 AND CHLORINE.

These were conducted into cooled silicon tetrachloride, as in the former instance. There was no liberation of heat, the liquid becoming merely colored by the gases and giving them up freely on distillation, the boiling-point remaining normal, thus indicating that no compound had been formed and that simply an absorption of the vapors had taken place. This is all the more remarkable, since the derivatives $3TiCl_4.2NO_2Cl$ and $3SnCl_4.2NO_2Cl$ had been readily prepared by Hampe.[3]

THE CHLORIDES OF PHOSPHORUS.

Phosphorus trichloride, under ordinary conditions, or heated under pressure with silicon tetrachloride, gives no derivative; the liquids merely commingle to a solution from which both constituents may be isolated by fractional distillation. Phosphorus in solid condition remains insoluble in silicon tetrachloride, but when its vapors, partly dissociated, are conducted into the same, they are absorbed and dissolved. Upon subsequent distillation

[1] *Ann. Chem.* (Liebig), **38**, 17.
[2] *Ann. Chem.* (Liebig), **39**, 320.
[3] *Ann. Chem.* (Liebig), **126**, 43.

the liquid, for the greater part, boils at 58°–60°, leaving a liquid residue, which is driven over only at the boiling-point of phosphorus trichloride. The only effect of the phosphorus pentachloride seems to be a slight elevation of the boiling-point of the silicon compound, while the latter serves to completely dissociate the pentachloride into the lower chloride and free chlorine. Casselmann[1] had prepared a compound of tin and phosphorus chloride, of the formula $(SnCl_4)_2.PCl_5$. Here again the analogy between tin and silicon fails.

SULPHUR MONOCHLORIDE.

The liquids were mixed in all proportions, and on boiling the mixture, the silicon tetrachloride being in excess, the greater part came over between 57°–59°. The temperature rises gradually and then suddenly is elevated to 139°, the boiling-point of sulphur monochloride, thus showing that no compound had been formed. H. Rose,[2] however, obtained the compound $SnCl_4.2S_2Cl_2$.

CYANOGEN CHLORIDE.

A saturated solution of mercuric cyanide was made, and an excess of the salt was added to this in a finely divided condition. Chlorine was then conducted into this liquid, until it was completely saturated by the gas, and the product was set aside in a dark place in a flask, filling the space above the liquid with chlorine. After a time the excess of mercuric cyanide had dissolved and the chlorine was completely absorbed. Uncombined chloride was then removed by shaking with mercury, and the solution was poured into a flask provided with a calcium chloride tube. On gently heating the solution, the cyanogen was evolved, and led through a tube filled with copper turnings, to remove the last traces of chlorine, into silicon tetrachloride chilled in ice and salt. Silicon in its behavior here was found to ally itself with tin, both Wöhler[3] and Klein[4] having failed to obtain chlorcyanogen compounds with the higher chlorides of the latter, while Wöhler had however succeeded in preparing a titanium derivative of the composition $TiCl_4(CNCl)_2$.

[1] *Ann. Chem.* (Liebig), **83**, 258.
[2] Pogg. *Annalen*, **42**, 517.
[3] *Ann. Chem.* (Liebig), **74**, 86.
[4] *Ann. Chem.* (Liebig), **73**, 221.

SILICON TETRACHLORIDE AND AROMATIC AMINES.

Knop[1] obtained derivatives from silicon tetrafluoride with urea and aniline. With the latter a white crystalline compound was prepared, to which he assigned the formula $Si_2H_3Fl_9.3Cl_2H_7N$. This constitutes the first work investigating the action of the aromatic amines on the tetrahaloids of the fourth group. As, however, the medium here used is anhydrous alcohol which decomposes the silicon tetrafluoride, the resulting compound cannot be said to be a derivative of it. Laurent and Delbos[2] obtained a compound of aniline and silicon tetrafluoride, whose constitution they expressed as $3C_6H_5NH_2.2SiF_4$, but did not establish. Jackson and Comey[3] obtained also this compound, proved its constitution, and described its properties. They prepared, in a similar manner, derivatives from ortho- and para-toluidine, diphenylamine and dibenzylamine, while dimethyl-aniline and quinoline gave no product.

Girard and Pabst[4] investigated the action of the tetrachlorides of carbon, tin, and silicon on aniline at increased temperatures. With carbon tetrachloride and aniline, they obtained the derivatives, triphenylguanidine and rosaniline. With the chloride of tin under the same conditions, violaniline, mauvaniline (phenyl-safranine), and it seems, rosaniline were obtained. Silicon tetrachloride also forms violaniline, together with the production of triphenylamine blue.

The only investigations on these reactions which are established by exact analyses are those of Harden[5] and Reynolds[6]; the latter is a repetition of the work of the former experimenter, in which results are obtained which vary from it. Unaware of any conflict of results, or, indeed, of any precedence in the field, the present investigation was begun; and it was not until its results were complete that any divergence in previous work on the subject was encountered. The work was then found to agree entirely with that of the earlier of these investigators, and to establish by new methods of procedure his results.

1 *Chem. Centrbl.*, *1858*, 388; *Jsb. d. Chem.*, *1858*, 148; *J. prakt. Chem.*, **74**, 41.
2 *Ann. chim. phys.*, **22**, 101.
3 *Ber. d. chem. Ges.*, **19**, 3194.
4 *Bull. Soc. Chim.*, **34**, 38.
5 *J. Chem. Soc.*, **51**, 40.
6 *J. Chem. Soc.*, **55**, 474.

SILICON TETRACHLORIDE AND ANILINE.

When aniline is added to silicon tetrachloride, there is a great evolution of heat, with the coincident precipitation of a white compound completely insoluble in ether. Owing to the violence of the reaction, anhydrous ether, and afterwards benzene, was used as a diluent and medium of interaction. The complete combination of the silicon tetrachloride was evidenced by the loss of the odor of that body when sufficient aniline had been added. With this as a means of indication, the synthetic proportions were found to be one molecule of silicon tetrachloride to four molecules of aniline.

Three and five-tenths grams silicon tetrachloride required 7.50 grams aniline. Calculated, 7.65 grams aniline.

Five and eight-tenths grams silicon tetrachloride required 12.4 grams aniline. Calculated, 12.68 grams aniline.

That the silicon tetrachloride united with the aniline in the proportion of one to four molecules, was further proved by weighing the resultants of the action of quantities of these substances united according to the above ratio. Ether was used as a medium.

To three and five-tenths grams of silicon tetrachloride, dissolved in ether, a slight excess of four molecules of aniline was added, stirring vigorously during the addition of the latter substance. The white precipitate thus obtained was placed on a tared filter and washed with anhydrous ether, in a current of dry air, until the washings showed no trace of aniline. The precipitate was then dried in a desiccator, the filter and precipitate placed in a weighing bottle and its weight found.

From three and five-tenths grams silicon tetrachloride were obtained 11.16 grams product. Calculated for three and five-tenths grams silicon tetrachloride, and 7.66 grams, or four molecules of aniline, gave 11.3 grams product.

In a second experiment the following figures were obtained:

Four and six-tenths grams of silicon tetrachloride gave 14.8 grams product. Calculated for 4.60 grams silicon tetrachloride and 10.06 grams, or four molecules of aniline, 14.66 grams product.

The additional weight of the precipitate was here probably due

to some small quantities of aniline retained by it. Portions of this substance which had been placed in the cup of a Soxhlet apparatus and washed with ether for three hours, were taken for analysis. The exit tube of the Soxhlet apparatus was, in this instance, connected with a drying apparatus, and the precipitate during the washing thus protected from the action of moist air. All precautions were taken to remove every trace of free aniline, the washings of the precipitate being constantly examined for the same, till they failed to give reactions of that body. The compound was placed in a desiccator and portions taken for analysis.

I. 0.1135 gram gave 0.0122 gram silica, or 5.052 per cent. silicon.

II. 0.3739 gram gave 0.0280 gram silica, or 4.80 per cent. silicon.

III. 0.0965 gram gave 0.0103 gram silica, or 5.01 per cent. silicon.

IV. 0.1432 gram gave 0.2805 gram carbon dioxide, or 53.42 per cent. carbon; and 0.0684 gram water, or 5.31 per cent. hydrogen.

V. 0.2153 gram gave 0.1572 gram Pt, or 10.47 per cent. nitrogen.

VI. 0.2034 gram gave by Dumas' method 0.02166 gram nitrogen, or 10.65 per cent. nitrogen.

VII. 0.3467 gram gave 0.3710 gram silver chloride, or 26.46 per cent. chlorine.

VIII. 0.2972 gram gave 0.3130 gram silver chloride, or 26.05 per cent. chlorine.

	Calculated for $(C_4H_8NH_2)_4SiCl_4$.	Found. I.	II.	III.
Silicon	5.16	5.05	4.80	5.01
Carbon	53.13	53.42	....	....
Hydrogen	5.16	5.31	....	....
Nitrogen	10.32	10.47	10.65	....
Chlorine	26.19	....	26.46	26.05

These analyses further substantiate the proportions of combination which had been indicated by the synthetic experiments. As the compounds thus produced possess the formula

$SiCl_4(C_6H_5NH_2)_4$, the reaction may be written as follows:

$$SiCl_4 + 4C_6H_5NH_2 = SiCl_4(C_6H_5NH_2)_4.$$

This was regarded as a single molecular compound, and the reaction was thus written until the use of a medium other than benzene, in the combination of these bodies, gave results that pointed to the conclusion that the above compound was a mixture of the compounds $SiCl_2(C_6H_5NH)_2$ and $2C_6H_5NH_2HCl$, so formed by the combination of one molecule of silicon tetrachloride and four of aniline. The resulting dichlorosilicon diphenylamide and the aniline hydrochloride, being both insoluble in ether, would be combined in such proportions as to give a mixture whose total composition would correspond to the formula deduced from the above analysis:

$$SiCl_2(C_6H_5NH)_2 + 2C_6H_5NH_2HCl = SiCl_4(C_6H_5NH_2)_4.$$

The results that led to the inference that the latter compound was not simple are below detailed. When instead of ether as a medium, anhydrous benzene is used, all of the products are not precipitated, but it was found on washing the precipitate thus obtained with benzene in a dry Soxhlet apparatus, a white compound was left, which gave the following proportions on analysis:

I. 0.3401 gram gave 0.3751 gram silver chloride, or 26.99 per cent. chlorine.

II. 0.2266 gram gave 0.2506 gram silver chloride, or 27.07 per cent. chlorine.

The compound thus obtained did not show any silica content.

	Calculated for aniline hydrochloride.	Found. I.	II.
Chlorine	27.13	26.99	27.07

It was further noticed that the total weight of this precipitate was only about half that calculated for the complete precipitation of the compound of silica, expressed in the reaction with ether as a medium.

I. Three and five-tenths grams silicon tetrachloride and four molecules of aniline gave 5.29 grams of product insoluble in benzene; with ether as a medium obtained 11.3 grams of product insoluble in ether.

II. Three and nine-tenths grams silicon tetrachloride gave 5.06 grams of product insoluble in benzene; calculated for $(C_6H_5NH_2)_4SiCl_4$, 12.43 grams.

As the aniline hydrochloride was entirely free from silica, the silicon compound produced in the reaction was looked for in the clear filtrate from the aniline hydrochloride. This solution was not allowed to evaporate, to yield in this manner the dissolved derivative, as it was known, since the aniline was originally in slight excess, that any compound so produced would be saturated with aniline, the last traces of which it would be difficult to remove. To the solution was therefore added ether, which precipitated a white compound containing silicon, carbon, hydrogen, nitrogen, and chlorine, and whose analyses gave the following results:

I. 0.2461 gram substance gave 0.4602 gram carbon dioxide, or 50.95 per cent. carbon; and 0.0860 gram water, or 3.88 per cent. hydrogen.

II. 0.2840 gram gave 0.0589 gram silica, or 9.69 per cent. silicon.

III. 0.2131 gram gave 0.0442 gram silica, or 9.74 per cent. silicon.

IV. 0.2761 gram gave 0.2788 gram silver chloride, or 24.70 per cent. chlorine.

V. 0.3013 gram gave 0.2014 gram Pt, by method of Varrentrapp and Will, or 9.59 per cent. nitrogen.

	Calculated for $SiCl_2(C_6H_5NH)_2$.	Found. I.	II.
Carbon	50.88	50.95	....
Hydrogen	4.24	3.88	....
Nitrogen	9.87	9.59	....
Silicon	9.87	9.69	9.74
Chlorine	25.08	....	24.70

This is the compound which had been obtained by Harden.[1] On weighing the amount of this produced from definite amounts of silicon tetrachloride, and adding to it the quantity of aniline hydrochloride produced at the same time, the total weight both of the silicon tetrachloride and the aniline are united and found in the products described above.

[1] *J. Chem. Soc.*, 51, 40.

Three and five-tenths grams of silicon tetrachloride and four molecules, or 7.65 grams, aniline gave :

	Grams.
Dichlorosilicon diphenylamide	5.83
Aniline hydrochloride	5.29
Total found	11.12
Calculated for three and five-tenths grams silicon tetrachloride and 7.65 aniline	11.15

Three and nine-tenths grams silicon tetrachloride and 8.55 grams aniline gave :

	Grams.
Dichlorosilicon diphenylamide	6.32
Aniline hydrochloride	5.96
Total found	12.58
Calculated for three and nine-tenths grams silicon tetrachloride and 8.53 grams aniline	12.43

These bodies are almost quantitatively produced, as the above results have indicated, and no amount of any secondary products can be believed to be present. The reaction may, therefore, be written as follows :

$$SiCl_4 + 4C_6H_5NH_2 = SiCl_2(C_6H_5NH)_2 + 2C_6H_5NH_2HCl.$$

Both these products were obtained, and only these, although the aniline was always in excess ; while Reynolds writes the reaction thus :

$$SiCl_4 + 8C_6H_5NH_2 = SiCl_2(C_6H_5NH)_2 + 2C_6H_5NH_2HCl.$$

The silicon tetraphenylamide prepared by him is a well-crystallized body, and the careful methods of analysis pursued place its exact constitution beyond doubt. The cause of the divergence in the results is, therefore, due to the variation in procedure. In the work of Hardin in the present investigation, smaller quantities of material are used (only a slight excess of four molecules of aniline are employed), and the diluents of the action were not present in such great amount as in the work of Reynolds, where the benzene is double the value of the reagents. In his work, 100 grams of silicon tetrachloride are added to 438 grams of aniline, or in the proportion of one to eight molecules.

In the use of such a quantity of benzene, the purpose is to keep the larger amount of the silicon compound in the solution, so that the precipitate will consist almost wholly of aniline hydrochloride, retaining but little of the silicon derivative. The precipitate, according to the method of this investigator, is quickly filtered from the solution, washed often with benzene and the filtrate distilled free from the greater part of the benzene in a current of hydrogen. The sirupy liquid thus obtained was poured into an excess of carbon disulphide, which served to precipitate a little aniline hydrochloride which had continued in solution. The filtrate was then concentrated and allowed to crystallize, when the compound described by him was obtained.

It will be here observed that the work is less direct than that of the methods of procedure before described, and that more chance of secondary reactions is encountered in the manipulation than in the first instance, and that the reaction which requires four molecules of aniline with one of silicon tetrachloride, represents the normal behavior of these two substances.

It is believed that the use of ether in lieu of benzene, furnishes a far preferable medium for determining the proportions of the reaction, though preventing an analysis of the two products, which are equally insoluble in it. It renders possible, however, a weighing of the total products of the reaction and a comparison of their quantity, with the amount of silicon tetrachloride and aniline used, since it exerts no solvent action on either product, while both reagents are soluble in it. The possibility of secondary reactions, or of the solvent playing the rôle of a reagent was thus avoided. The readiness of weighing the total product of the reaction, and the determination of the synthetic proportions of the same, seem to establish that it proceeds as follows :

$$4C_6H_5NH_2+SiCl_4 = SiCl_2(C_6H_5NH)_2+2C_6H_5NH_2HCl,$$

and that the compound without chlorine, $Si(C_6H_5NH)_2$, and the reaction which Reynolds writes, does not represent the immediate action of the reagents, but that the time-consuming method of the latter introduces a second phase into the reaction, during which the following may take place :

$$SiCl_2(C_6H_5NH)_2+4C_6H_5NH_2 = Si(C_6H_5NH)_4+2C_6H_5NH_2HCl,$$

which would explain the formation of both silicon derivatives and the contradiction of the reactions.

It might here be noticed that the names "Dichlorosilicon diphenylamide" for $SiCl_2(C_6H_5NH)_2$, and "Silicon tetraphenylamide" for $Si(C_6H_5NH)_4$, are cumbrous and not sufficiently clear; "Chlorsilicon dianilide," for the first, and "Silicon tetranilide" for the latter, are suggested as briefer and more expressive names.

SILICON TETRACHLORIDE AND ORTHOTOLUIDINE.

The methods employed in the preparation of the toluidine derivatives were similar to those made use of in the production of the aniline compound. A greater reactivity was, however, noticed on the part of the toluidine, as it was found to combine with the silicon tetrachloride at the temperature of ice and salt, while aniline was inert under these conditions. The precaution was taken to remove all paratoluidine from the liquid variety, by means of their acetyl derivatives, and the pure orthotoluidine thus formed was used in the reactions.

The white compound insoluble in benzene was washed with ether, dried, and analyzed.

I. 0.3146 gram gave 0.3126 gram silver chloride, or 24.32 per cent. chlorine.

II. 0.4012 gram gave 0.4006 gram silver chloride, or 24.43 per cent. chlorine.

Calculated for $C_6H_4CH_3NH_2HCl$, 24.73 per cent. chlorine.

The compound is, therefore, toluidine hydrochloride and its production similar to that of the aniline body.

On the addition of ether to the filtrate, a white voluminous precipitate appeared, which was washed with anhydrous ether, and dried in a desiccator. Of this substance:

I. 0.1722 gram gave 0.0319 gram silica, or 8.70 per cent. silicon.

II. 0.1650 gram gave 0.0313 gram silica, or 8.91 per cent. silicon.

III. 0.2160 gram gave 0.0403 gram silica, or 8.77 per cent. silicon.

IV. 0.1834 gram gave 0.1727 gram silver chloride, or 22.51 per cent. chlorine.

V. 0.2803 gram gave 0.2617 gram silver chloride, or 22.85 per cent. chlorine.

	Calculated for $SiCl_2(C_6H_4CH_3NH)_2$.	Found. I.	II.	III.
Silicon	9.00	8.70	8.91	8.77
Chlorine	22.80	22.51	22.85	....

The behavior of orthotoluidine is, therefore, analogous to that of aniline. That four molecules of toluidine were required for the saturation of silicon tetrachloride was proved as in the former instance by determining the synthetic proportions of the compounds, and by the use of ether as a medium, determining the weight of the total product. It required :

	Toluidine. Grams.
For 3.0 grams silicon tetrachloride	7.7
Calculated	7.6
For 5.3 grams silicon tetrachloride	13.2
Calculated	13.2

The precipitate produced and washed in ether in the

	Product. Grams.
First experiment gave	10.8
Calculated for 7.6 and 3.0	10.6
Second experiment gave	18.8
Calculated for 5.3 and 13.2	18.5

SILICON TETRACHLORIDE AND ACID AMIDES. BENZAMIDE.

The vapors of silicon tetrachloride, conducted into benzamide in the molten condition, gave rise to benzaldehyde, which was collected and recognized by the malachite green reaction. It was thought that the silicon tetrachloride would form a product with the benzamide, as with the aromatic amines, but it is probable that the benzamide is not sufficiently basic in character to so act.

SILICON TETRACHLORIDE AND ACETAMIDE.

With the latter compound no derivative is obtained, but on conducting the vapors into acetamide, acetonitrile was produced. Possibly according to the following :

$$2CH_3CONH_2 + SiCl_4 = 2CH_3CN + SiO_2 + 4HCl$$

The present investigation on the behavior of silicon tetrachloride is but one in a series which has for its scope the determination of the reactions also of lead and tin tetrachlorides with amine bodies. Work is also in progress in this laboratory on the conduct of thorium and zirconium tetrachlorides with the same reagents. The results thus obtained will afford a wider study of the fourth group elements, and furnish data for a broader comparison of the reactions of its members.

SOME NEW RUTHENOCYANIDES AND THE DOUBLE FERROCYANIDE OF BARIUM AND POTASSIUM.

By Jas. Lewis Howe and H. D. Campbell.[1]

Received November 26, 1897

THE resemblance between the double cyanides of iron, ruthenium, and osmium, was first pointed out by Claus,[2] and further developed by Martius,[3] who formed a number of the double osmocyanides. More recently Dufet[4] has described the crystals of potassium ruthenocyanide and osmocyanide and shown their close crystallographic resemblance to the ferrocyanide. The chemistry and the crystallography of the ferrocyanides has been very exhaustively studied by Wyrouboff,[5] and his work affords a valuable starting-point for any investigation of the isomorphous ruthenocyanides and osmocyanides.

Since the publication of a recent paper by one of us (Howe)[6] on the ruthenocyanides, several new salts have been prepared, and in all of them the analogy of the ruthenocyanides with the ferrocyanides is complete. The ruthenocyanides are decidedly more soluble than the ferrocyanides and correspondingly more difficult to crystallize. In one salt only have crystals been obtained which were perfect enough to measure.

Strontium Ruthenocyanide, $Sr_2Ru(CN)_6.15H_2O$.—Formed by treating lead ruthenocyanide with dilute sulphuric acid and neutralizing of the hydroruthenocyanic acid formed with strontium

[1] Contributed to the November meeting of the Cincinnati Section of the American Chemical Society, November 16, 1897.

[2] Beiträge zur Chemie der Platinmetalle. pp. 98 and ff.

[3] Ueber die Cyanverbindungen der Platinmetalle. Inaug. Diss., Göttingen, 1860.

[4] *Compt. rend.*, 120, 377.

[5] *Ann. chim. phys.* [4], 16, 280; 21, 271.

[6] This Journal, 18, 981.

hydroxide. The salt obtained by treating "Prussian purple," the ruthenium analogue of Prussian blue, with strontium hydroxide, could not be obtained in a pure condition. Pale straw-colored, elongated plates, probably monoclinic, very soluble in water, and quite efflorescent. By adding alcohol to its hot solution, the salt can be obtained in long, fine needles. It is very difficult to obtain a salt pure enough for analysis, and owing to efflorescence the water determination is low. The corresponding strontium ferrocyanide crystallizes with fifteen molecules of water of crystallization.[1]

ANALYSIS.

	I. Crystallized from water. Calculated for $Sr_2Ru(CN)_6, 15H_2O$.	Calculated for $Sr_2Ru(CN)_6.14H_2O$.	II. Crystallized from dilute alcohol. Found. I.	II.
Water	38.41	36.79	36.80	37.12
Ruthenium	14.45	14.83	14.36	....
Strontium	24.94	25.59	25.11	....

Barium Potassium Ruthenocyanide, $K_2BaRu(CN)_6, 3H_2O$.—Formed by mixing equivalent solutions of the barium and potassium salts; also mixed with the barium salt, by treating Prussian purple with barium hydroxide. Crystallizes best from its hot solution in water while evaporating on the water-bath. Pale amber to colorless rhombohedra, hardly distinguishable from the corresponding ferrocyanide, but quite soluble in water. The salt is difficult of analysis, owing to the difficulty of completely decomposing by heat to obtain the insoluble oxide of ruthenium. Some of the ruthenium invariably went into solution in hydrochloric acid, and from this solution a blue precipitate, insoluble in acids, soon separated, probably $Ru(CN)_2$, but invariably containing both barium and potassium.

ANALYSIS.

	Calculated for $K_2BaRu(CN)_6, 3H_2O$.	Found. I.	II.	III.	IV.
Potassium	14.83	12.98	....	14.51	13.16
Barium	26.07	27.38	25.31	....	25.83
Ruthenium	19.27	18.71	...	....	18.94
Water	10.25	10.19	10.80	10.83	10.99

The water in I was determined by loss, but in the other three analyses by absorption and direct weighing. In these three

[1] Wyrouboff: *Ann. chim. phys.*, 16, 287.

cases the salt in a closed tube was heated until it began to decompose. It is therefore impossible that the salt can contain five molecules of water, as this would require sixteen per cent.

Wyrouboff[1] describes the barium potassium ferrocyanide as $K_2BaFe(CN)_6,5H_2O$, but crystallographically the two salts resemble each other very closely.

Crystallography of barium potassium ruthenocyanide, $K_2BaRu(CN)_6,3H_2O$. Rhombohedral. $a : c = 1.558$. Angle between basal plane and rhombohedron $= 119° 4'$. The mean of thirty-nine measurements on eight crystals gave $81° 27'$ as the rhombohedral angle. (Calculated $= 81° 36'$.)

Crystals rhombohedral, with occasional development of basal plane. Rhombohedral faces usually appear pitted, and give a bundle of reflections, making a variation of $50'$ in the measured angles; optically negative; double refraction weak. In converging polarized light the cross opens slightly into hyperbolas.

A comparison of the crystallography of the ruthenocyanide with the ferrocyanide as given by Wyrouboff shows their close relationship.

Barium potassium ferrocyanide.[2]	Barium potassium ruthenocyanide.
Rhombohedral. $a : c = 1.570$.	Rhombohedral. $a : c = 1.558$.
Angle between basal plane and rhombohedron $= 118° 53'$.	Angle between basal plane and rhombohedron $= 119° 4'$.
Rhombohedral angle $= 98° 48'$ ($81° 12'$).	Rhombohedral angle $= 81° 36'$ (calculated); $81° 27'$ (measured).
Optically positive.	Optically negative.

Barium Cesium Ruthenocyanide, $Cs_2BaRu(CN)_6,3H_2O$.—When barium ruthenocyanide is mixed with half the equivalent of cesium sulphate, there separate from the solution, even before it can be filtered from the barium sulphate, unless very dilute, minute, brilliant, yellowish-white to colorless, crystals, which under the microscope appear as rhombohedra, with angle of about 83°, and in every way resembling the barium potassium salt, except that the basal plane predominates. They are very insoluble even in hot water, and cannot be recrystallized owing to partial decomposition and precipitation of a blue insoluble

[1] Wyrouboff: *Ann. chim. phys.*, 21, 279.
[2] Wyrouboff: *loc. cit.*

substance, probably $Ru(CN)_2$. The salt is stable in the air as analyses II and III were made from a salt which had stood exposed in the laboratory for several weeks.

ANALYSIS.

	Calculated for $Cs_2BaRu(CN)_6,3H_2O$.	Found. I.	II.	III.
Cesium	37.18	....	....	36.45
Barium	19.21	19.06	....	18.70
Ruthenium	14.21	....	....	14.33
Water (by loss at 120°)	7.56	7.64	7.31	7.51

The sodium, the magnesium, and the calcium salts have been obtained, but are very difficult to crystallize, owing to their great solubility. The magnesium salt forms very pale green crystals which are very efflorescent.

The strontium potassium, and the strontium cesium ruthenocyanide have been formed, but not in pure enough condition for analysis. They seem to resemble the barium double salts under the microscope and bear no resemblance to the salt $K_2SrFe(CN)_6$, $3H_2O$, described by Wyrouboff, which crystallizes in prismatic monoclinic crystals. The strontium rubidium ruthenocyanide gives a precipitate on warming its solution, and thus appears to resemble the strontium potassium ferrocyanide which precipitates the anhydrous salt under similar circumstances.

Analysis having shown that the barium potassium ruthenocyanide contains three molecules of water of crystallization, and yet resembles closely the ferrocyanide to which Wyrouboff[1] gives five molecules, a reexamination of the ferrocyanide was made. Bunsen, who first made the salt, gives the formula with three molecules of water, and from analogy to the ruthenocyanide three would be expected. Further, Martius[2] formed the corresponding osmocyanide, to which he gives the formula $K_2BaOs(CN)_6,3H_2O$. His description of the salt would hold equally good for the ruthenocyanide or ferrocyanide, except that it is efflorescent.

A considerable quantity of the ferrocyanide was prepared in this laboratory by Mr. H. W. Drake. The very sparingly soluble barium salt was made by the action of barium chloride on potassium ferrocyanide, and this, mixed with an equivalent

[1] Wyrouboff: *loc. cit.*

[2] Martius: Cyanverbindungen, p. 23.

quantity of the potassium salt was dissolved in a large amount of water. On evaporation and cooling the barium potassium salt was deposited in pale straw-colored rhombohedra, often several millimeters across. The salt, contrary to the statement of Wyrouboff, is perfectly stable, and after five months' exposure to the air the crystals are perfectly clear and the surfaces bright. Measurement of the crystal angles, and examination of the optical properties, showed the salt to be identical with that described by Wyrouboff. Analysis showed that the salt has *three* molecules of water of crystallization, and its formula is $K_2BaFe(CN)_6,3H_2O$. The water was determined directly by absorption in a calcium-chloride tube, the salt being heated until decomposed.

ANALYSIS.

	Calculated for $K_2BaFe(CN)_6,5H_2O$.	Calculated for $K_2BaFe(CN)_6,3H_2O$.	Found. I.	Found. II.
Water	17.40	11.38	11.68	11.72
Barium	26.55	28.53	28.88	28.61

Barium Cesium Ferrocyanide, $Cs_2BaFe(CN)_6,3H_2O$. This salt, which appears never to have been described, was made for purposes of comparison. It was formed by decomposing barium ferrocyanide with half the equivalent of cesium sulphate in very dilute solution. It crystallizes in very small, slightly yellowish-white rhombohedra, and is very difficultly soluble in water. Its hot solution appears to be stable.

ANALYSIS.

	Calculated for $Cs_2BaFe(CN)_6,3H_2O$.	Found.
Cesium	39.71	38.03
Barium	20.53	20.42
Iron	8.37	8.53
Water	8.07	8.01

Crystallography, rhombohedral. $a : c = 1.515$. Angle between basal plane and rhombohedron $= 119° 45'$ (calculated). Rhombohedral angle $= 82° 30'$.

This salt adds confirmatory testimony to the amount of water of crystallization in barium potassium ferrocyanide, and also to the close crystallographical resemblance between the salts of the type $M'BaM'''(CN)_6,3H_2O$. ($M' = K_2Cs_2$; $M''' = Fe, Ru, Os$).

WASHINGTON AND LEE UNIVERSITY,
LEXINGTON, VA., Nov. 13, 1897.

THE CAFFEIN COMPOUND IN KOLA. PART II. KOLATANNIN.[1]

By James W. T. Knox and Albert B. Prescott.

THE investigation of the caffein compound in kola conducted by us last year[2] showed us that it was a kolatannate of caffein instead of a glucoside as formerly supposed. This question then arose: Is the tannin of this body identical with the free tannin existing in the kola nut? In the very brief examination made at that time it seemed that there was some difference, for combustions of both kinds of the tannin showed a difference in the respective amounts of carbon and hydrogen of each. No interpretation of the results was made at that time, for we considered the data insufficient, and the question was not settled.

It became necessary, in order that this point might be positively determined, to make a thorough comparative study of these two forms of kolatannin, and the research has been continued along this line with results which are stated in the following pages.

CLASSIFICATION OF TANNINS.

Several different means of classifying tannins have been proposed, such as "physiological" and "pathological"—referring, of course, to their origin in the plant; glucosidal and non-glucosidal; "iron-blueing" and "iron-greening;" tannins which are tanning agents and those which are not; those which yield a trihydroxyphenol by sublimation or by fusion with potassium hydroxide, and those which yield a dihydroxyphenol by sublimation, or the salt of a dihydroxybenzoic acid by gentle fusion with potassium hydroxide.[3]

Trimble[4] suggests a classification, not as a final one but which answers very well for the present, in view of the still very defective knowledge of this class of bodies. He proposes a division of them into "the gall tannin group" and "the oak tannin group." The tannins of the gall tannin group contain 50 to 52.5 per cent. of carbon and 3.10 to 4.50 per cent. of hydrogen; those of the oak tannin group contain, in round numbers, sixty

[1] Presented by the Special Research Committee at the Forty-fifth Annual Meeting of the American Pharmaceutical Association, August, 1897.

[2] Knox and Prescott 1896: This Journal, 19, 63.

[3] Prescott: Organic Analysis, p. 466.

[4] Trimble: The Tannins, II, p. 131.

per cent. carbon and five per cent. hydrogen. This classification holds good for the following reactions also :

	Gall tannin group.	Oak tannin group.
Ferric salts.	Blue color and precipitate	Green color and precipitate.
Calcium hydroxide	White precipitate becoming blue.	Light pink precipitate becoming red and then brown.
Bromine water.	No precipitate.	Yellow precipitate becoming brown.

Under this classification kolatannin is to be compared with the oak tannin group, for its compositions and reactions coincide more nearly with those of oak tannin than with those of gallotannic acid. We include herewith a brief summary of some important work done on oak bark tannins, not only for its historic interest, but because it has a bearing indirectly upon our own work on kolatannin.

RECENT LITERATURE OF OAK BARK TANNINS.

In February, 1880, Böttinger[1] published a lengthy article on "oak red," phlobaphene and tannin, which was followed about a month later by Etti's[2] publication of his own work on oak tannin—a valuable contribution to its literature. The methods used by them in the preparation of their working samples differed radically, the botanical source of their barks or extracts was unknown to them (or, if known, was not stated), the tannins obtained possessed properties different from each other, and it is not strange that the results obtained by these investigators were at variance in many particulars. If we may judge by their descriptions, it would seem that Etti's method of separation was much more likely to yield a pure product than that used by Böttinger, although by no means free from objections. Without doubt this work of Etti and that of Löwe,[3] dated in September of the same year, but which did not appear until the following year, added more to the knowledge of oak tannins than any previous contributions had done. The methods used by them are noteworthy, because they were radical departures from the old lead acetate methods formerly so much in vogue.

[1] C. Böttinger, 1880: *Ann. Chem.* (Liebig), **202**, 269.
[2] C. Etti, 1880: *Monatsh. Chem.*, **1**, 262.
[3] J. Löwe, 1881: *Ztschr. anal. Chem.*, **20**, 208.

Etti extracted the oak bark with very dilute alcohol under gentle heat and added acetic ether and ordinary ether to the extract. After agitating, the faintly red-colored ethereal layer was separated, the ether recovered by distillation and used again for the same purpose. After distilling the ether, the alcoholic solution of tannin remaining behind was found to contain some crystals of ellagic acid, which were removed by filtration and the filtrate evaporated to dryness on a water-bath. The residue, a reddish-white powder, was oak tannin, mixed with a little phlobaphene, amorphous resin and gallic acid; the two last-named substances were separated by extracting the powder with non-alcoholic ether; it was next treated with a mixture of three parts acetic ether and one part ethyl ether, which removed the tannin and left the phlobaphene undissolved. After distilling off the solvent, the pure tannin remained behind as a reddish-white powder. Etti insists, too, that if the ethyl acetate be used again for a solvent it must be washed with sodium hydroxide solution and redistilled before use, as the tannin tends to decompose it slightly into acetic acid and alcohol, while the tannin is itself partially converted into its anhydride.

The method of Böttinger depends on the separation of the phlobaphene from tannin solutions by means of its comparative insolubility in water. The tan-bark is extracted first with ether, then with alcohol, the alcohol solution is evaporated to dryness, and the residue extracted again with ether which removes the last traces of wax and fat. By warming the residue with water on a water-bath, a partial separation is effected of the insoluble phlobaphene from the soluble tannin. After filtration the tannin solution, still containing traces of phlobaphene, is repeatedly evaporated and redissolved, until upon diluting the liquid very largely and cooling it, no precipitation is had. On evaporating this solution, a light-colored residue is obtained which is soluble in water and which he considered the pure tannin.

Löwe extracted the oak bark with ninety per cent. alcohol, and distilled the extract *in vacuo* to sirupy consistency. Water was then added, which precipitated a large quantity of anhydride; after filtration common salt was added to saturation, which precipitated the remaining anhdyrides completely; the liquid, after filtration, was agitated with ether repeatedly until this sol-

vent would remove nothing further from the solution. The remaining ether was dissipated by gentle heat and after cooling the solution was shaken out with acetic ether. The acetic ether was removed by distillation and cold water was added to the remaining residue as long as precipitation ensued, after which it was again filtered and evaporated slowly in a vacuum desiccator to a dry mass, which is soluble in water and becomes cinnamon-brown on being pulverized.

It will be noticed that the methods of Etti and Löwe have much in common, the principal difference from the other methods being in the use of ethyl acetate, an immiscible solvent. It is difficult to say to whom credit should be given for the first application of this solvent to the preparation of tannin. In 1873,[1] however, Löwe mentions having used it in his work on sumach tannin, while in 1872[2] he also made use of it as a separative solvent for gallotannic acid.

Etti ascribed the formula $C_{17}H_{16}O_9$ to his tannin: a "first anhydride" $C_{34}H_{30}O_{17}$ was formed by heating the tannin at 140° C. He separated a ready-formed anhydride from the bark corresponding to this one, and made use of its barium salt $C_{34}H_{28}BaO_{17}$ to determine its formula. By boiling it with dilute hydrochloric or sulphuric acid, it loses another molecule of water, forming a second anhydride $C_{34}H_{28}O_{16}$. The third anhydride he prepared by boiling the pure tannin with dilute mineral acid, whereby two molecules of tannin lost three molecules of water, forming a body to which he ascribed the formula $C_{32}H_{26}O_{15}$. This anhydride corresponds, in Etti's opinion, to Oser's oak-red. Boiling the pure tannin with potassium hydroxide solution for some time, formed the first anhydride identical with the so-called oak bark phlobaphene; this was separated by acidulating the alkaline solution with mineral acid and filtering out the precipitated anhydride.

By various experiments, such as boiling the tannin with dilute acids, digesting it with ferments, etc., he arrived at the conclusion that oak tannin was not a glucoside but that such evidence of glucose as had been found by other investigators was due to the accidental presence of glucose in the sample.

He proposed a structural formula, the same as that of digallic

[1] J. Löwe, 1873: *Ztschr. anal. Chem.*, **12**, 128.
[2] J. Löwe, 1872: *Ztschr. anal. Chem.*, **11**, 365.

acid with three hydroxyls groups replaced by three methyl groups, he having detected methyl chloride by burning the gas formed by heating the tannin with concentrated hydrochloric acid in a sealed tube.

Böttinger's work was principally on oak phlobaphene and oak-red, which he considered identical. The oak-red was formed by boiling the tannin with dilute mineral acid. He ascribed the formula $(C_{14}H_{10}O_6)_2.H_2O$ to it and believed oak tannin to be a glucoside.

Löwe proposed the formula $C_{28}H_{28}O_{14}$ for the tannin he prepared and $C_{28}H_{22}Pb_2O_{14}$ for the lead salt. He also prepared oak-red by the usual process of boiling the tannin with dilute acids and obtained in addition to the oak-red, $C_{28}H_{22}O_{11}$, a by-product of which he made a combustion, but did not identify. It was not sugar, however, and he states that oak tannin does not conduct itself as a glucoside. The oak phlobaphene is, according to his statements, a fourth anhydride of oak tannin of the formula $C_{28}H_{24}O_{12}$, and forms a lead salt $C_{28}H_{22}O_{12}Pb$.

A somewhat extended and not altogether good-natured controversy between Böttinger and Etti followed the publication of the latter's paper, Böttinger[1] insisting upon the correctness of his own work, and holding to his first statement that oak tannin was a glucoside, Etti,[2] on the other hand, contended that Böttinger's methods were faulty, that ready-formed glucose would not be separated from the tannin by Böttinger's process, and therefore sugar would of course be found in the solution after boiling the tannin with dilute acids. He performed a number of experiments tending to prove the correctness of his position, following Böttinger's methods also and criticising him severely. Böttinger[3] subsequently modified his position somewhat, but without conceding anything of importance.

In 1883 the next contribution appeared, in which Etti[4] again commented on Böttinger's work and quoted Löwe in confirmation of his own results. He reported a new tannin, of formula $C_{20}H_{20}O_9$, from *Quercus Pubescens*, while the other he had since

1 C. Böttinger, 1881 : *Ber. d. chem. Ges.*, **14**, 1598.
2 C. Etti, 1881 : *Ber. d. chem. Ges.*, **14**, 1826.
3 C. Böttinger, 1881 : *Ber. d. chem. Ges.*, **41**, 2390.
4 C. Etti, 1883 : *Monatshefte*, **4**, 512.

learned was from *Quercus Robur*. This new tannin colored solutions of ferric salts green, while the first one produced a blue color with them. During this investigation the objection already noted to the use of ethyl acetate, *viz.*, its tendency to decompose the tannin, became so marked as to necessitate a different method of separation. The most important points of this new method are the use of alcoholic ether as a solvent for the tannin, phlobaphene and green resin. The resin is removed from the resulting product by benzene, and the phlobaphene by careful addition of lead acetate to the tannin solution. The precipitate of lead tannate is yellow, but if any anhydride (phlobaphene) be present, the precipitate will be colored more or less reddish-brown, depending on the proportion of it to the tannin. Lead acetate is added until the color of the precipitate indicates that all of the anhydride has been precipitated. After filtering, the liquid is again exhausted with alcoholic ether, the ether removed by distillation and the residual liquid evaporated to dryness on the water-bath. A rather remarkable peculiarity of Etti's work is that both of these tannins are nearly insoluble in water, 0.6 of tannin in 100 of water, while the experience of other workers has been quite generally that tannins are soluble.

Etti claimed to have formed four anhydrides of this tannin, $C_{20}H_{20}O_9$.

First anhydride,	$C_{40}H_{38}O_{17}$.	Oak phlobaphene, dried at 120°.
Second "	$C_{40}H_{36}O_{16}$.	By boiling phlobaphene with 1 : 12 sulphuric acid and washing precipitate first with water, then with alcohol, and evaporating the alcoholic solution.
Third "	$C_{40}H_{34}O_{15}$.	Dried at 125°. By boiling tannin with dilute sulphuric acid and collecting the precipitate.
Fourth "	$C_{40}H_{32}O_{14}$.	By heating in 25 per cent. sulphuric acid containing 20 per cent. alcohol in sealed tube at 130° for four hours.

Etti believed also that four anhydrides were formed from the other tannin ($C_{17}H_{16}O_9$) and gave this view support by his own figures and some also of Oser, of Böttinger and Löwe. This would be apt to mislead the casual reader into attaching more importance to this theory of the four anhydrides, than is war-

ranted by the actual results given by Etti. It would hardly seem that he was justified in making this use of the work of the other investigators. As a matter of fact he has selected figures here and there which gave support to his views, but ignored others of equal importance which did not harmonize with his ideas; *e. g.*, he quoted Löwe's figures for the analysis of oak-red, but did not mention his work with the lead salts of oak-red, which brought him to the conclusion that the formula of oak-red was $C_{28}H_{22}O_{11}$. There is no apparent reason for doubting that this work was equally as reliable as that which Etti quoted, and to omit it gives a careful reader the impression that deductions made from such peculiarly isolated facts would not necessarily be of great value. The chart is here appended:

		Calculated.	Found.			
Tannin ..... $C_{17}H_{16}O_9$	Carbon	56.04	56.31	56.06		
	Hydrogen ..	4.40	4.63	4.69		
First anhydride ..... "Phlobaphene," .... $C_{34}H_{30}O_{17}$ ..... Grabowski's oak-red.	Carbon	57.46	57.28	57.62		
	Hydrogen ..	4.22	4.64	4.33		
Second anhydride ... $C_{34}H_{28}O_{16}$	Carbon	58.96	58.76			
	Hydrogen ..	4.04	4.20			
			By Oser.	By Böttinger.	By Etti.	
Third anhydride .... $C_{34}H_{26}O_{16}$ ..... Oser's oak-red.....	Carbon	60.58	60.70	60.19	60.08	60.33
	Hydrogen ..	3.86	4.03	4.22	3.94	4.03
			Found by Löwe.			
Fourth anhydride ... $C_{34}H_{24}O_{14}$ ..... Löwe's oak-red.....	Carbon	62.20	62.339	62.197	61.997	
	Hydrogen ..	3.66	4.154	4.015	4.056	

Böttinger[1] published in the same year, 1883, a piece of work done on the bromine derivatives of oak tannin. By direct addition of bromine to the watery extract of oak bark he formed a dibrom oak tannin, $C_{19}H_{14}Br_2O_{10}$, which was capable of forming a pentacetyl derivative. "By careful manipulation"—he does not describe it—he converted the dibrom into a tetrabromtannin, which also formed a pentacetyl compound. This seems to be the first work on bromine derivatives.

In 1884[2] he investigated hemlock tannins by means of their

[1] C. Böttinger, 1883: *Ber. d. chem. Ges.*, **16**, 2710.

[2] C. Böttinger, 1884: *Ber. d. chem. Ges.*, **17**, 1041.

bromine compounds precipitated directly from their infusions by addition of bromine. Later[1] in the same year he prepared bromine compounds of a number of dark tannins by adding bromine directly to their infusions. Acetyl compounds of these were formed and analyzed.

Etti,[2] in a brief paper shortly afterward, discussed the points of difference between gallotannic acid and oak tannin. He also pointed out that by reason of its very hygroscopicity acetic anhydride was not a proper agent for determining the number of hydroxyl groups of tannins, as its tendency would be to form anhydrides which in turn would be acetylized and lead to incongruous results in the analysis.

Böttinger,[3] in 1887, published an article principally on oakwood tannin, which he separated by dissolving commercial oakwood extract in twenty parts of water and allowing it to stand. The clear liquid was evaporated to dryness and boiled with acetic anhydride, which formed an acetyl compound whose composition corresponded to formula $C_{15}H_7(C_2H_2O)_5O_9$. From this he recovered an anhydride of oakwood tannin and formed mono- and tetrabrom derivatives of it.

In 1889, Etti,[4] adhering to his previously published formulas for oak tannins, gave the results of an investigation of a tannin from the common Slavonian oak, to which he gave the formula $C_{16}H_{14}O_9$. It was nearly or quite insoluble in water. He formed several soluble magnesium salts with this tannin and suggested that it might exist in the plant in this form, thus accounting for its solubility in an aqueous menstruum when first extracted. Several anhydrides were formed.

Any résumé of the literature of oak tannins, however brief, would be incomplete without reference to the work of Trimble.[5] His monograph in two volumes contains a praiseworthy compilation of previous literature on the subject and full accounts of his own work. He has the credit of introducing acetone as an extractive menstruum for the oak tannins instead of alcohol. This solvent is said to act particularly well with oak bark. A

[1] C. Böttinger, 1884: *Ber. d. chem. Ges.*, **17**, 1123.
[2] C. Etti, 1884: *Ibid*, **17**, 1820.
[3] C. Böttinger, 1887: *Ann. Chem.* (Liebig), **238**, 366.
[4] C. Etti, 1889: *Monatshefte*, **10**, 547.
[5] H. Trimble, 1894: "The Tannins." A monograph, J. B. Lippincott Co.

good bibliography concludes the work. He has confined himself thus far to analyses of the pure tannins and does not as yet propose formulas. In addition to the monograph just mentioned, he has been a frequent contributor on this subject in the periodical literature of chemistry.

LATE WORK OF SCHIFF ON DIGALLIC ACID.

In the present year, Hugo Schiff,[1] having already found that natural tannic acid is possessed of a slight optical activity, namely, dextrorotatory, inquires as to its containing in some way asymmetric carbon, and infers a ketonic structure with six hydroxyl groups. He does not obtain a hydrazone from tannic acid. From experimental results he proposes this formula:



Irrespective of orientation, and for comparison of a kolatannin formula offered for consideration further on, the above formula may be held simply as

$$\begin{matrix} C_6H(OH)_3(CO_2H) \\ C_6H_2(OH)_2 \end{matrix} \Big\rangle CO.$$

CAFFEOTANNIC AND CAFFEIC ACIDS.

The literature of caffeotannic acid, of the formula $C_{15}H_{18}O_8$, dates from Rochleder, in 1846.[2] Its production of caffeic acid (dihydroxyphenylacrylic acid), along with sugar, was set forth by Hlasiwetz in 1867.[3]

A few years ago, caffeic acid was obtained from the tannin of coffee, also from the tannin of the maté, by H. Kunz-Krause.[4] Very recently the same author has investigated caffeotannic

1 *Gazz. chim. ital.*, *1897*, **27**, 1, 90.
2 *Ann. Chem.* (Liebig), **59**, 300.
3 *Ibid.*, **142**, 220.
4 *Arch. Pharm.*, *1893*, **231**, 613.

acid.[1] In his report he[2] finds confirmation of the composition of caffeic acid as a dihydroxyphenylacrylic acid. His analyses obtain the carbon dioxide given off when the caffeic acid is heated to 200° C., and are confined to this analytic method. A residue of a vinyl pyrocatechol is predicted. The author recognizes a class of bodies which he terms glucotannoids.

"FREE" KOLATANNIN.

PREPARATION OF THE SAMPLE.

The amorphous character of tannins, together with their extreme liability to decomposition, renders their separation and purification very difficult. The method used by us last year,[3] in the very brief examination of kolatannin, was based on that of Allen,[4] being fractional precipitation with lead acetate.

On taking up the work this year, it soon became evident that whatever may be said of lead-acetate methods as applied to other tannins, they certainly are unsuited to this one, for various reasons. The lead kolatannate, being bulky, is extremely slow and difficult of filtration, and the tannin in this state exhibits a marked tendency to pass into insoluble forms, principally anhydrides. During the washing, the lead salt becomes gradually darker, and this tendency is favored by the moisture present. If, on the other hand, the washing be less prolonged, the resulting product will be contaminated with various other constituents of kola extract, such as sugar, alkaloids, inorganic salts, coloring-matter, etc., etc. It is indeed very doubtful if any usual amount of washing will remove these impurities, hence the purity of a tannin, separated by precipitation with lead acetate, would be regarded with too much suspicion to be employed in analytical work, even if its sensible properties were not changed during the treatment. Kolatannin was at first supposed and since determined to be soluble in water. But on evaporating the solution of tannin after removing the lead from combination by hydrogen sulphide, the end-product was a red-brown

[1] I am indebted to Prof Kremers for bringing Kunz-Krause's article in the July *Berichte* to my attention, as I read this paper in the August meeting of the American Pharmaceutical Association.—A. B. P.

[2] *Ber. d. chem. Ges.*, *1897*, 1617.

[3] Knox and Prescott, *loc. cit.*

[4] Allen: Com. Org. Anal., 3, pt. I., 76.

substance, insoluble in water, and possessing a not particularly astringent taste. So by the use of lead acetate a tannin is obtained whose purity is a matter of doubt, and whose properties have undergone radical changes during separation, both as to color and solubility. Furthermore, we desired to effect a perfect separation of the free tannin from that existing in combination with caffein (as caffein kolatannate) in order to determine by analysis whether they were identical tannins or not, the analysis of them last year having shown somewhat different values in each for carbon and hydrogen. Caffein kolatannate is sparingly soluble in water, more soluble in solutions of tannin and of caffein, so that the aqueous infusion of kola contains caffein kolatannate as well as free tannin, and the lead tannate obtained from it would be a mixture of the lead salts of both tannins, or both forms of tannin, and no delicate separation could be had in this way.

Plainly, another method was necessary. The objections just noted were for the most part neither new, nor limited to kolatannin. As previously stated, Löwe, in 1872, who had encountered many difficulties in the use of lead acetate for the separation of tannins, and found it necessary to employ other means to obtain a pure product, resorted to the use of ethyl acetate. This immiscible solvent, proving satisfactory in many cases, has since been used considerably, either alone or mixed with ether, in the separation of tannins for analytical purposes, and for some tannins at any rate is unquestionably the best separative solvent which has yet been given publicity as such. At the same time a single shaking out with ethyl acetate does not suffice for the preparation of a pure sample. More or less coloring-matter will be found in the residue left on evaporating the first portion of the solvent, so that several repetitions of the process are required. Care must be used not to carry the method of purification too far, for if this be done the decomposition of the tannin is apt to prove a greater obstacle than the original impurities. We find that four or five repetitions of the "shaking out" gave the best results. In detail the manipulations, are as follows, and the method has given satisfactory results :

The fresh kola nuts are sliced into boiling alcohol,[1] removed after a few moments boiling, and dried in a current of warm air, then ground up to a number twenty powder and packed firmly in a percolator. The alcohol so used in sterilizing the drug is diluted to about fifty per cent. strength, and employed as a menstruum, with addition of sufficient dilute alcohol to complete the extraction. The highly-colored extract of kola thus obtained is concentrated by distillation *in vacuo*, until the alcohol is all removed. This can be done with a good pump at between 18° and 20° C. The contents of the flask are then filtered, the insoluble portion being chiefly caffein kolatannate, while the solution contains caffein, kolatannin, caffein kolatannate, glucose, traces of fatty matter, and more or less colored derivatives of tannin according to whether the sterilization by boiling alcohol has been properly done. Common salt is then added to this filtrate to saturation and the caffein kolatannate is completely precipitated. It is filtered out and added to the first residue of caffein kolatannate. The reddish-colored filtrate is then transferred to a large separatory funnel and agitated with chloroform repeatedly to remove alkaloids and traces of fat. The dissolved chloroform is then separated by agitation with small portions of ether. Kolatannin is insoluble in chloroform, and only very sparingly soluble in ether, so this treatment does not remove any considerable amount of it. Ethyl acetate is now added to the liquid in the separatory funnel and the liquid extracted with it as long as any tannin is removed. Usually eight, ten, or even more successive portions are required for this. The united solutions of tannin in ethyl acetate are transferred to a distilling flask and concentrated to dryness under reduced pressure. To avoid loss of the ethyl acetate we use a condenser about a meter in length, and pack the receiver in ice. Sometimes also a condenser is placed between the pump and the receiver. With these precautions the waste is very slight. Before using the recovered ethyl acetate as a solvent again, however, it is washed with a weak solution of sodium carbonate and redistilled, to avoid the presence of any free acetic acid formed by decomposition of the ether. The statement of Etti already alluded to, concern-

[1] To prevent the formation of the colored body, which would otherwise appear.

ing the decomposition of ethyl acetate by tannin, has not been found to apply to kolatannin, but the washing was nevertheless adopted as a precautionary measure.

The tannin residue in the distilling flask is a porous, pinkish mass, very friable, and easily and completely soluble in water. It is redissolved in a convenient quantity of saturated salt solution, filtered if need be and again shaken out with ethyl acetate, which is distilled off in the way just described. The tannin residue is next dissolved in cold distilled water and shaken out as before, this process being repeated once or twice. The final tannin residue, after the ethyl acetate has been removed as far as possible by distillation, is treated in the flask with a small quantity of ether, which after thoroughly permeating the mass is removed by distillation, using only the partial force of the pump and gentle heat. This is repeated several times until the mass has no odor of ethyl acetate. Then transferring to a vacuum desiccator the tannin is kept for several days over sulphuric acid. If this does not remove the last traces of ether, gentle heat (60°–65°) will do so.[1]

PROPERTIES OF KOLATANNIN.

Kolatannin is a cream-colored powder with a slight pinkish tinge. It is freely and completely soluble in water, alcohol, acetone, ethyl acetate, sparingly soluble in ether, insoluble in chloroform and in petroleum benzine. The following tabulated comparative statement of its reactions will serve to point out some of the important differences between it and gallotannic acid, and to indicate the close resemblance between this tannin and that of oak bark.

[1] We desire to take this means of expressing our appreciation of the liberality and courtesy of the firm of Frederick Stearns & Company, Manufacturing Pharmacists Detroit, Michigan, who supplied us with a large quantity of True African Kola nuts, for our research this year. The drug was quite fresh and in perfect preservation, very necessary conditions for work of this kind.

REACTIONS OF TANNINS.

	Kolatannin. Free	Kolatannin. From caffein kolatannate.	Oak tannin. (Trimble.)	Gallotannic acid.
Ferric acetate	Green.	Green.	Green.	Blue-black.
Potassium dichromate	Dark brown ppt.	Dark brown ppt.	Brown ppt.	Brown ppt.
Chlorine	Pale ppt.	Pale ppt.		
Bromine	Light yellow ppt.	Light yellow ppt.	Yellow ppt.	No ppt.
Calcium hydroxide	Pink, then red, changing to ppt.	Pink, then red, changing to ppt.	Red, changing to ppt.	Darkens with bluish tinge, then ppt.
"Tartar Emetic"	No ppt.	No ppt.	White ppt.	White ppt.
Quinin ... Cinchonin ... Caffein ...	White ppt.	White ppt.	Ppt.	White ppt.
Albumen	Ppt.	Ppt.	Ppt.	Ppt.
Lead acetate	White ppt.	White ppt.	White ppt.	White ppt.
Ammoniacal potassium ferricyanide	Deep Red.	Deep red.	.....	Deep red.
Solution of formaldehyde with a condensing agent, as HCl	Pink ppt., becoming red.	Pink ppt., becoming red.		

COMBUSTIONS OF KOLATANNIN.

The tannin was analyzed by combustion with the following stated results;

	Found.			Calculated for
	I.	II.	III.	$C_{14}H_{10}O_8$.
Carbon	56.74	56.81	56.90	56.45
Hydrogen	5.71	5.62	5.76	5.90
Oxygen	37.55	37.57	37.34	37.65
	100.00	100.00	100.00	100.00

Having this satisfactory evidence of the preparation of a pure chemical compound, we next proceeded upon the task of determining its constitution, first by finding the number of hydroxyl groups in the molecule containing C_{14}.

PENTACETYL KOLATANNIN.

An acetyl derivative was next formed for the object above stated. The tannin was boiled with acetyl chloride for about one hour under a return condenser, then an equal volume of glacial acetic acid was added, after which the liquid was poured slowly into ice-cold water. The acetyl compound separated out as a voluminous whitish precipitate. This was rapidly filtered at the pump and well washed with cold water, then it was dried with bibulous paper and finally in a vacuum desiccator over sulphuric acid. It was a nearly white or gray-white powder, tasteless, having a faint odor of acetic acid, and was insoluble in water, sparingly soluble in ether, and freely and completely soluble in chloroform, in alcohol, and in glacial acetic acid.

Analysis of the body gave the following stated results:

	Found.			Calculated for
By combustion.	I.	II.	III.	$C_{14}H_{18}(C_2H_3O)_8O_8$.
Carbon	56.55	56.68	56.76	56.72
Hydrogen	5.39	5.36	5.24	5.47
Oxygen	38.06	37.96	38.00	37.81
By saponification.				
Acetyl	38.24	38.47	38.68	39.09

On comparing the values given by analysis with those calculated for the pentacetyl derivative, the agreement seems to be satisfactory.

The acetyl values in the analysis by saponification as detailed below are a little low, but it is quite possible that this is to be

ascribed to defects in the method of estimation rather than to other causes. At any rate the acetyl content seems without doubt to indicate five hydroxyl groups in the C_{14} molecule rather than four or six.

The analyses by combustion were made in the ordinary way: the estimation of acetyl was conducted in the manner next described.

DETERMINATION OF ACETYL.

The acetyl compound, one-half to one gram, varying with the supposed acetyl content, is boiled with thirty to forty cc. of six per cent. sulphuric acid in a flask fitted with reflux condenser, for two hours. 100 to 150 cc. water are now added and without filtration the volatile contents of the flask distilled into a known quantity of decinormal potassium hydroxide solution, being careful to avoid a high enough temperature or sufficient concentration to cause distillation of the sulphuric acid. More water is now added to the distilling flask and the distillation repeated. The contents of the receiver should be kept alkaline and a few drops of phenolphthalein solution added, so that any change in the reaction of the solution may be instantly noted. The distillation should be repeated several times until the last one hundred cc. of distillate contains no acetic acid. The excess of alkali is to be titrated with decinormal hydrochloric acid, and the amount of acetic acid calculated from the difference between the amount of alkali solution first taken, and the quantity of hydrochloric acid required to neutralize it. Barium chloride is now added, and if any precipitation takes place, the barium sulphate is collected and weighed, the sulphuric acid calculated and correction made for it. If a halogen be present in the acetyl compound, a sufficient excess of silver sulphate is added to the contents of the distilling flask before distillation, to precipitate it completely as silver salt.

The above was occasionally deviated from by heating the acetyl compound with the dilute acid in sealed tubes two or three hours at 130°, then cooling, opening, transferring to a flask and distilling as before described.

The following described method was used a few times, but without apparent advantage over the others: The sample is

boiled with twenty times its weight of alcoholic solution of potassium hydroxide, of five to eight per cent. alkali strength, in a flask under return condenser. Phosphoric acid is added in excess and the contents of the flask distilled with steam into a receiver containing a known quantity of decinormal alkali solution. To drive the acetic acid over completely requires distillation of about a liter of water. The same precautions are observed as in the preceding methods if halogens are present.

BROMINE DERIVATIVES OF KOLATANNIN.

The bromine compounds of this tannin have been investigated at considerable length, for the purpose of obtaining further data concerning its constitution.

TRIBROM KOLATANNIN.

When bromine water is added to a water solution of kolatannin, a dark-colored body is precipitated, and with continued addition of the reagent the precipitation goes on until complete, during which the precipitate formed becomes lighter-colored until at the end, in the presence of an excess of bromine, it is pale brownish yellow and further addition of bromine water produces no change. It is quite voluminous and cannot well be filtered at the pump, but is collected by ordinary filtration and well washed with water until the washings remain clear when treated with silver nitrate solution. It is dried first on porous plates, then in a vacuum desiccator to constant weight.

During the process of drying it gradually changes color until it becomes reddish brown. It is oderless and nearly tasteless, insoluble in water, ether, chloroform, and benzene, but is readily soluble in alcohol and in acetone.

In the clear filtrate, after the bromine compound has been removed, hydrobromic acid is easily identified, and for that reason the change is believed to be one of substitution. It may be questioned perhaps whether this is not caused by the action of bromine on the water; the bromine water, however, was freshly prepared, and the hydrobromic acid formed under these conditions would be very little. But in preparing twenty-five or thirty grams of this bromine compound, after its precipitation, the filtrate was agitated with carbon disulphide to remove the excess

of bromine, then separated and the aqueous portion concentrated to small volume, which had an acid taste and reaction and in which hydrobromic acid was easily identified.

The bromine compound was analyzed with the following stated results:

	Found.			Calculated for
By combustion.	I.	II.	III.	$C_{18}H_{17}Br_3O_8$.
Carbon	32.91	33.12	....	33.28
Hydrogen	2.69	2.58	....	2.95
Oxygen	22.86	22.63	....	22.19
As silver salt.				
Bromine	41.54	41.67	41.62	41.58

The percentage of carbon in this body, when compared with that of the pure tannin from which this was prepared, seemed to indicate the substitution of three hydrogen atoms by that number of bromine atoms.

PENTACETYL TRIBROM KOLATANNIN.

(From Tribrom Kolataunin.)

For the purpose of obtaining further data, an acetyl compound of the bromine derivative was formed: The bromine compound is boiled with acetyl chloride for about forty-five minutes under a reflux condenser. An equal volume of glacial acetic acid is added to the clear red solution and the liquid well cooled by freezing-mixture. Then it is poured slowly into twelve or fifteen times its volume of ice-cold water, which is constantly stirred. A thermometer should be kept in the beaker containing the precipitate, and if the temperature rises above 10°, it is best to cool the contents by placing the beaker in a freezing-mixture, resuming addition of the acetyl chloride on cooling, until all has been used. The acetyl compound separates as a flocculent yellow precipitate, is quickly filtered, well washed and dried, first on porous plates, then in a vacuum desiccator.

It is a tasteless, golden-yellow powder, having a faint odor of acetic acid. It is insoluble in water, nearly insoluble in ether, but soluble in alcohol, in acetone, and in chloroform. It loses acetic acid when gently heated, and both bromine and acetic acid on stronger heating. It was analyzed with the following stated results:

	Found.		Calculated for
By combustion.	I.	II.	$C_{16}H_{12}Br_2.(C_2H_3O)_6O_8$.
Carbon	39.43	39.55	39.65
Hydrogen	3.44	3.34	3.44
Oxygen	26.57	26.65	26.43
As silver salt.			
Bromine	30.56	30.46	30.48
By saponification.			
Acetyl	26.91	27.12	27.32

The results obtained agree very closely with the calculated formula and give additional confirmation to the formulas assumed for the pure tannin and its tribrom compound.

HALOGEN DETERMINATION.

For the halogen determinations in the bromine compounds of kolatannin, the well-known process of heating the halogen derivative with fuming nitric acid and silver nitrate in a sealed tube for two hours at 175°, was followed. Experience taught us, however, to first make a preliminary determination with a considerable excess of silver nitrate to ascertain the approximate amount of halogen present. For example, where in one case the preliminary determination indicated 40.88 per cent. of bromine, the following ones, in triplicate with only a slight excess of silver nitrate, were respectively 41.54 per cent., 41.67 per cent., 41.62 per cent. And as a rule the preliminary determinations have given low results, but are necessary to furnish data for the second and more exact estimations.

Having learned the approximate amount of halogen present, the following details of manipulation were followed: The size of the sample taken is adjusted so that the weight of the resulting silver salt shall be 0.15 to 0.20 gram. Six or eight per cent. more than the theoretical quantity of silver nitrate is now weighed and placed with the sample in a hard glass tube with six cc. fuming nitric acid, after which the tube is sealed and heated at 175°–180° for two hours. It is then opened and the contents are transferred to a beaker with about 100 cc. water and heated on the water-bath for one-half hour. The silver bromide is then collected on weighed asbestos filters connected with the pump, washed successively with hot water, alcohol and ether, and dried twenty or thirty minutes at 80°–85° to constant weight.

In some cases the excess of silver nitrate in the filtrate was deter-

mined gravimetrically as silver chloride for a check on the results.

Our thanks are due to Professor Edward D. Campbell, of this University, for many valuable suggestions on halogen and acetyl determinations.

TRIBROM PENTACETYL KOLATANNIN.

(From Pentacetyl Kolatannin.)

Having acetylized the tribrom compound of tannin and obtained a pentacetyl derivative of it, we decided to brominate the pentacetyl compound and determine by analysis whether or not the resulting product was identical, in other words to ascertain whether the order in which the substitutions were made affected the result.

This bromination was accomplished by dissolving the pentacetyl tannin in strictly pure, anhydrous chloroform, free from alcohol, and adding a slight excess of bromine. The solvent was then distilled off in partial vacuum, while the brominated acetyl compound remained behind as a golden-yellow mass. This was treated with ether several times and redistilled, using the pump to remove the traces of chloroform, and finally transferred to a vacuum desiccator and left until completely dry.

In color, taste, and solubilities this body is identical with that obtained by acetylizing the tribrom derivative. It was analyzed with the following stated results:

	Found.		Calculated for
By combustion.	I.	II.	$C_{18}H_{19}Br_3(C_2H_3O)_5O_8$.
Carbon	39.49	39.62	39.65
Hydrogen	3.39	3.47	3.44
Oxygen	26.61	26.34	26.43
As silver salt.			
Bromine	30.51	30.57	30.48
By saponification.			
Acetyl	26.84	26.98	27.32

Inasmuch as the results of the analysis of this body coincide quite as closely with the theoretical figures as in the preceding case, it may be safely assumed that the bodies are identical and that in the preparation of a pentacetyl tribrom kolatannin it makes no difference whether the bromine or acetyl is introduced first.

TETRABROM KOLATANNIN.

We next wished to ascertain whether this tannin was capable of taking more than three bromine atoms into combination in each molecule. Nothing but the tribrom derivative was formed by treating an aqueous solution of the tannin with bromine water in excess, so another solvent was tried successfully. Taking a weighed quantity of tannin, it was dissolved in alcohol and bromine was added gradually to the alcoholic solution in slight excess of the theoretical quantity necessary to form a tetrabrom derivative. The liquid remained clear, all the bromine compounds of kolatannin that we have yet prepared being easily soluble in alcohol. The bromine compound was precipitated by pouring the brominated alcoholic solution into twelve or fifteen volumes of ice-cold water: being insoluble in water, it was separated as a voluminous reddish brown precipitate, which was filtered, washed and dried in the way already described under the method of preparation of tribrom kolatannin.

This body agrees in solubilities with the tribrom compound; it is slightly darker, however, and has a faint odor of bromine. It was analyzed with the following stated results:

	Found.		Calculated for
By combustion.	I.	II.	$C_{18}H_{16}Br_4O_8$.
Carbon	29.59	29.46	29.27
Hydrogen	2.46	2.38	2.44
Oxygen	18.97	18.88	19.52
As silver salt.			
Bromine	48.98	49.28	48.77

This, then, is a tetrabrom derivative of kolatannin. From it an acetyl compound was formed, following the manipulations already described.

PENTACETYL TETRABROM KOLATANNIN.

This acetyl compound was yellow and but slightly darker than that of the tribrom derivative. In other physical properties it appeared identical with that body.

It was analyzed with the following stated results:

By combustion.	Found. I.	II.	Calculated for $C_{16}H_{11}Br_4(C_2H_3O)_5O_8$.
Carbon	35.82	35.97	36.03
Hydrogen	3.02	2.95	3.01
Oxygen	24.02	24.14	24.02
As silver salt.			
Bromine	37.14	36.94	36.94
By saponification.			
Acetyl	24.42	24.59	24.83

From the above it is seen that the tetrabrom derivative forms a pentacetyl compound equally as easily as the tribrom derivative.

PENTABROM KOLATANNIN.

This derivative was formed while trying to prepare a hexabrom substitution product, but by reason of an unsufficiency of bromine the resulting product yielded inalytical results corresponding fairly well with those calculated for pentabrom kolatannin.

By combustion.	Found. I.	II.	Calculated for $C_{16}H_{16}Br_5O_8$
Carbon	25.87	25.99	26.13
Hydrogen	2.07	1.86	2.04
Oxygen	17.35	17.31	17.42
As silver salt.			
Bromine	54.71	54.84	54.41

This body may be a mixture of the hexa- and tetrabrom tannins. It is somewhat difficult of formation, however, and we have not always been able to obtain it in a manner analogous to the methods of preparation of the other bromine compounds. We do not consider that all conditions necessary to its formation are yet understood. There is no good evidence that this is a mixture of the other derivatives, however, and for the present there is really no sufficient reason for mentioning it as other than a pentabrom tannin. It is soluble in the same agents as the other bromine compounds, and is less stable than the tetrabrom compound, the odor of bromine being much more distinct. It appears to be characteristic of these bodies, that with increasing percentage of bromine, there is decreasing stability (that is to say, a greater tendency to lose bromine), and darker color.

PENTACETYL PENTABROM KOLATANNIN.

(From Pentabrom Kolatannin.)

This was prepared by acetylzinig pentabrom kolatannin in the usual way. In properties and appearance it closely resembles the other bromacetyl derivatives, except that its color is somewhat darker yellow. It was analyzed with the following stated results:

	Found.		Calculated for
By combustion.	I.	II.	$C_{16}H_{10}Br_5(C_2H_3O)_5O_8$.
Carbon	33.54	33.39	33.02
Hydrogen	2.77	2.65	2.65
Oxygen	20.97	21.37	22.01
As silver salt.			
Bromine	42.72	42.59	42.32
By saponification.			
Acetyl	22.15	22.36	22.75

This body was also prepared from pentacetyl kolatannin, as stated next below.

PENTABROM PENTACETYL KOLATANNIN.

(From Pentacetyl Kolatannin.)

Having ascertained by previous experiment that addition of a considerable excess of bromide to an alcoholic solution of kolatannin, resulted in the formation of a hexabrom derivative, we wished to prepare a hexabrom pentacetyl derivative by brominating the pentacetyl derivative in the same manner.

Analysis of the resulting product showed, however, that contrary to our expectations, not a hexabrom, but a pentabrom, derivative was formed, identical with the one described in the preceding section. The analytical results obtained were as follows:

	Found.		Calculated for
By combustion.	I.	II.	$C_{16}H_{10}Br_5(C_2H_3O)_5O_8$.
Carbon	33.19	32.97	33.02
Hydrogen	2.60	2.71	2.95
Oxygen	21.90	21.95	22.01
By silver salt.			
Bromine	42.31	42.37	42.32
By saponification.			
Acetyl	22.31	22.47	22.75

HEXABROM KOLATANNIN.

By treating an alcoholic solution of kolatannin with bromine

in considerable excess a hexabrom compound is formed, which is separated in the same way as the tetra- and pentabrom derivatives. It is darker colored than either of the others, has a very distinct odor of bromine, and is soluble in the same media as the other bromine compounds. It was analyzed with results stated next below:

	Found.		Calculated for
By combustion.	I.	II.	$C_{14}H_{14}Br_4O_8$.
Carbon	23.48	23.57	23.59
Hydrogen	1.68	1.79	1.72
Oxygen	15.73	15.38	15.73
As silver salt.			
Bromine	59.11	59.26	58.96

TETRACETYL HEXABROM KOLATANNIN.

(From Hexabrom Kolatannin.)

This acetyl compound was prepared in the usual way by boiling with acetyl chloride and was supposed, before analysis, to be a pentacetyl derivative. It had a dark yellow color, a faintly pungent taste and a slight odor of bromine and acetic acid. Soluble in chloroform and in alcohol, insoluble in water. The analytical results which are next stated show it to be a tetracetyl hexabrom tannin:

	Found.		Calculated for
By combustion.	I.	II.	$C_{14}H_{10}Br_6(C_2H_3O)_4O_8$.
Carbon	29.57	29.39	29.33
Hydrogen	2.21	2.32	2.25
Oxygen	19.41	19.21	19.55
As silver salt.			
Bromine	48.81	49.08	48.87
By saponification.			
Acetyl	16.86	17.19	17.52

INDICATIONS OF MOLECULAR CONSTITUTION.

From the analytical results of the pentacetyl pentabrom tannin and of the hexabrom compound with its tetracetyl derivative, it appears that the total number of groups or atoms which may be substituted in the way described is ten; that if five of acetyl be introduced first, only five bromine atoms may be substituted for hydrogen, but that if the bromine be introduced first, six bromine atoms can be introduced by carrying bromination to the limit, and in this hexabrom compound only four hydroxyl

groups remain. This appears to indicate that the sixth bromine atom introduced bears a relation to the molecule in some respect different from that of any other bromine atom. But it might be expected that a limit would be reached in the total number of negative groups or atoms which can be retained in the molecule. The greatest number of acetyl groups probably indicates the entire number of hydroxyls. And a balance of evidence seems to favor the conclusions that there are six atoms of hydrogen directly united with the carbon of a benzene chain. These numbers are all in ratio to sixteen atoms of carbon. That the molecule contains C_{16}, or some multiple of this, appears from the quantitative results. We attempted to find the molecular weight of kolatannin, tribrom kolatannin and pentacetyl tribrom kolatannin by the cryoscopic method, but the results so far obtained are utterly meaningless. Further indications of moloculer constitution may be drawn from a study of the anhydrides, account of which next follows, also from the results of the action of acids, and especially from the products by fusion with alkali, all stated further on in this paper.

THE ANHYDRIDES OF KOLATANNIN AND THEIR DERIVATIVES.

By judiciously heating kolatannin several anhydrides may be prepared, as described below, the amounts of water driven off varying with the temperature and the duration of its application. The first anhydride of the series, that in which two molecules of tannin lose one molecule of water, and whose empirical formula is $(C_{16}H_{18}O_7)_2O$, is formed by heating kolatannin at 107°–110° to constant weight. This anhydride is darker colored than the tannin from which it is prepared and is less soluble in water. The second anhydride, by an oversight, we did not attempt to prepare. If we may judge by the temperature required for the preparation of the first, third and fourth members of the series, this one would result from heating the tannin a sufficient time at 120°–125°. Its theoretical formula is $C_{16}H_{16}O_7$. The third anhydride, $(C_{16}H_{17}O_8)_2O$, results from heating the tannin at 135°–140°. It is dark reddish brown and nearly or quite insoluble in water. The fourth anhydride, $C_{16}H_{16}O_8$, is

dark brown, insoluble in water, and is the result of heating kolatannin two hours at 155°–160°.

These anhydrides are all soluble in alcohol, and in solution of kolatannin of sufficient concentration.

FIRST ANHYDRIDE.

The method of preparation has been described in the preceding section. This is a yellowish red body, soluble in water, in alcohol, and in dilute alkali solutions. The calculated loss of weight of the tannin in preparing this anhydride was 2.65 per cent. The observed loss of weight was 2.77 per cent. The body was subjected to combustion with the following stated results:

	Found. I.	Found. II.	Calculated for $(C_{14}H_{10}O_7)_2O$.
Carbon	58.27	58.39	58.18
Hydrogen	5.38	5.61	5.45
Oxygen	36.35	36.00	36.37

FIRST ANHYDRIDE OF TRIBROM KOLATANNIN.

The first anhydride of kolatannin was dissolved in water and precipitated by adding an excess of bromine, collecting the precipitate by filtration, and drying exactly as described in the method of preparation of tribrom kolatannin. It is soluble in the same media as tribrom kolatannin, is darker colored, is odorless and tasteless. Analysis of it proved to be an anhydride of tribrom kolatannin.

By combustion.	Found. I.	Found. II.	Calculated for $(C_{14}H_{14}Br_3O_7)_2O$.
Carbon	33.61	33.80	33.86
Hydrogen	2.92	2.76	2.65
Oxygen	21.31	21.17	21.16
As silver salt.			
Bromine	42.16	42.27	42.33

FIRST ANHYDRIDE OF TETRABROM KOLATANNIN.

Prepared in the same way as tetrabrom kolatannin, using in this case the first anhydride of kolatannin. It is hardly to be distinguished from the other tetrabrom compounds in appearance or other sensible properties. It was analyzed with the following stated results:

By combustion.	Found. I.	II.	Calculated for $(C_{16}H_{18}Br_4O_7)_2O$.
Carbon	29.47	29.66	29.72
Hydrogen	2.37	2.21	2.17
Oxygen	18.57	18.42	18.58
As silver salt.			
Bromine	49.59	49.71	49.53

FIRST ANHYDRIDE OF HEXABROM KOLATANNIN.

This was prepared by adding a considerable excess of bromine to the alcoholic solution of the first anhydride of kolatannin, and separating the product in the usual way. It is dark red, tasteless, and has a faint odor of bromine; is insoluble in water, ether, and chloroform; soluble in alcohol and in dilute alkali solutions. It was analyzed with the following stated results:

By combustion.	Found. I.	II.	Calculated for $(C_{16}H_{12}Br_6O_7)_2O$.
Carbon	23.96	24.15	23.88
Hydrogen	1.64	1.69	1.49
Oxygen	14.61	14.57	14.93
As silver salt.			
Bromine	59.79	59.59	59.70

THIRD ANHYDRIDE OF KOLATANNIN.

The method of preparation has been described. The calculated loss of weight on heating the tannin was 7.96 per cent.; the observed loss was 8.16 per cent. It is a red-brown powder, odorless, nearly tasteless, but somewhat astringent. Soluble in alcohol and in alkali solutions, insoluble in water, ether, and chloroform. It was analyzed with the following stated results:

By combustion.	Found. I.	II.	Calculated for $(C_{18}H_{17}O_8)_2O$.
Carbon	61.72	61.80	61.54
Hydrogen	5.29	5.46	5.13
Oxygen	32.99	32.74	33.33

The bromine derivative of this anhydride and those of the fourth anhydride to be described further on in these pages, were prepared in a manner strictly analogous to those already described.

THIRD ANHYDRIDE OF TETRABROM KOLATANNIN.

A dark-brown, tasteless powder having a faint odor of bromine.

Insoluble in water, ether, or chloroform; soluble in alcohol and in alkali solutions. The analytical results were:

By combustion.	Found. I.	II.	Calculated for $(C_{16}H_{12}Br_4O_6)_2O$
Carbon	30.59	30.73	30.57
Hydrogen	1.99	1.81	1.91
Oxygen	16.32	16.19	16.56
As silver salt.			
Bromine	51.10	51.27	50.96

THIRD ANHYDRIDE OF HEXABROM KOLATANNIN.

A dark-brown, tasteless powder having a distinct odor of bromine. It is insoluble in water, ether, and chloroform; soluble in alcohol and in alkali solutions.

The analytical results were:

By combustion.	Found. I.	II.	Calculated for $(C_{16}H_{11}Br_6O_6)_2O$.
Carbon	24.04	24.39	24.43
Hydrogen	1.52	1.38	1.27
Oxygen	13.22	12.88	13.23
As silver salt.			
Bromine	61.22	61.35	61.07

FOURTH ANHYDRIDE OF KOLATANNIN.

This was prepared in the way previously described. The calculated loss of weight attendant upon its preparation from kolatannin is 10.61 per cent. The observed loss is 10.78 per cent. It is a dark-brown, odorless, tasteless powder, soluble in alcohol and in alkali solutions; insoluble in water, in ether, and in chloroform. It was analyzed with the following stated results:

	Found. I.	II.	Calculated for $C_{16}H_{16}O_8$.
Carbon	63.20	62.98	63.37
Hydrogen	5.09	5.16	4.95
Oxygen	31.71	31.86	31.68

FOURTH ANHYDRIDE OF TETRABROM KOLATANNIN.

A dark-brown, tasteless powder having a faint odor of bromine. Soluble in alcohol and in alkali solutions; insoluble in water, in ether, and in chloroform. It was analyzed with the following stated results:

By combustion.	Found. I.	II.	Calculated for $C_{14}H_{12}Br_4O_6$.
Carbon	30.91	30.68	31.02
Hydrogen	1.05	1.86	1.78
Oxygen	15.40	15.58	15.51
As silver salt.			
Bromine	51.74	51.88	51.69

FOURTH ANHYDRIDE OF HEXABROM KOLATANNIN.

A very bark-brown powder having a distinct odor of bromine. Soluble in the same media as the previously described bromine compounds. The analytical results are:

By combustion.	Found. I.	II.	Calculated for $C_{14}H_{10}Br_6O_6$.
Carbon	24.96	24.79	24.71
Hydrogen	1.39	1.21	1.16
Oxygen	11.70	12.01	12.35
As silver salt.			
Bromine	61.95	61.89	61.78

PRODUCTS OF THE ACTION OF DILUTE ACIDS UPON KOLATANNIN.

When kolatannin is boiled with dilute sulphuric or dilute hydrochloric acid, it first passes into solution, and on continued boiling a reddish precipitate is formed, which becomes gradually darker the longer the heating is kept up. This precipitate is collected by filtration and well washed with water, in which it is insoluble. It is also insoluble in alcohol, and only partially soluble in alkali solution. Combustions of this substance, while giving fair duplicates for the same sample, indicates that its composition is not uniform and constant. Samples Nos. I and IV were prepared by boiling kolatannin with six per cent. sulphuric acid, in a flask fitted with a return condenser; No. II by heating the tannin and dilute acid in a sealed tube; No. III by boiling the tannin and dilute acid in a beaker with occasional addition of water to compensate for evaporation; No. V by boiling caffein tannate with dilute acid in a flask fitted with return condenser; No. VI is the result of combustion of a similar substance made and reported last year.[1]

	I. Av. of 2.	II. Av. of 2.	III. Av. of 2.	IV. Av of 2.	V. Av. of 2.	VI. Av. of 2.
Carbon	53.60	50.45	51.27	56.38	54.49	69.20
Hydrogen	5.34	4.85	5.24	5.69	5.87	6.70
Oxygen	41.06	44.70	43.49	37.93	39.64	24.10

[1] Knox and Prescott, 1896: This Journal, 19, 89.

This aqueous filtrate was shaken out with ether, the residue after evaporation of the ethereal solution was redissolved in water and boiled a short time, which caused precipitation of the coloring-matter. Then filtering, it was again extracted with ether, and the operation repeated several times. At length the etheral solution left a nearly colorless residue on evaporation, which responded to the tests for protocatechuic acid. Oak bark tannin also, when subjected to similar treatment with dilute mineral acids, yields protocatechuic acid.

Etti, Böttinger, and Löwe each mention the insoluble red body formed from oak tannin on boiling it with dilute acids and consider it an anhydride. They do not agree on which of the anhydrides it is, however. In fact, as already mentioned, there are many points in their work on tannins on which they were at variance. Etti[1] found gallic acid in the acid filtrate. Löwe[2] does not mention having examined the liquid for bodies other than sugar. Trimble,[3] however, whose work is the most recent, found protocatechuic acid as a by-product, in every case. He does not commit himself on the question of whether the dark brown, amorphous, insoluble substance, separated from tannin by the action of hot dilute acids, is an anhydride.

Whatever may be the relation of the body separated in this way from oak bark tannin to the original tannin, there does not as yet seem to us to be any good reason for considering the corresponding substance precipitated from acid solutions of kolatannin, an anhydride of kolatannin. The evidence thus far is against it. Boiling with dilute mineral acids is in general a means of hydrolysis, rather than anhydrolysis; we would therefore expect saponification instead of anhydride formation; as a matter of fact an aromatic acid—protocatechuic—has been found in the acid filtrate after collecting the insoluble red substance in question, while if the reaction were one of simple anhydride formation, no by-product should be present. If the red-brown substance were an anhydride of kolatannin, we would expect it to have a higher percentage of carbon than the original tannin, but in most cases combustion indicates a lower percentage of

[1] C. Etti, 1884: *Ber. d chem. Ges.*, 17, 1820.
[2] J. Löwe, 1881: *Ztschr. anal. Chem.*, 20, 208.
[3] H. Trimble: "The Tannins," II, p. 92.

carbon. We would also expect to find the body of uniform composition when prepared by similar means, but the analyses just mentioned show that it is very variable in content of carbon and hydrogen, while its appearance also varies from red to dark brown. At present no more can be said of this body than we have already said; *viz.*, that it is red or dark brown, amorphous, insoluble in water, in alcohol, in ether and in chloroform, only partially soluble in dilute alkali, and that it is a decomposition product obtained by boiling kolatannin, or caffein kolatannate, with dilute mineral acids.

Knebel's[1] "kolaroth" is a body obtained in the same way, except that he prepared it from caffein kolatannate, or as he supposed "kolannin," which was the alleged "caffein-bearing glucoside" of kola. Inasmuch as the free tannin of kola and that combined with caffein are identical, as will appear further on in these pages, it makes no difference which kind of it is used for this purpose. Knebel, however, worked with caffein tannate from the dried drug, while our work has been entirely on that from the fresh drug. He fails to give the figures of his combustion, but Hilger[2] reports the following stated results:

	Found.		Calculated for $C_{14}H_{12}(OH)_6$.
	I.	II.	
Carbon	63.09	62.65	63.16
Hydrogen	6.85	6.50	6.75
Oxygen	30.06	30.85	30.09

This calculated formula does not differ widely in carbon content from that calculated for the fourth anhydride of kolatannin, $C_{18}H_{16}O_8$.

	Calculated for $C_8H_{16}O_8$.
Carbon	63.37
Hydrogen	4.95
Oxygen	31.68

The hydrogen is rather high, however, which makes the oxygen correspondingly low. The results of an analysis of a body like the one Hilger analyzed might have been constructed to indicate it as a member of the anhydride series, and inasmuch as he found a substance yielding analytical figures nearly coinci-

[1] E. Knebel, 1892: *Apoth. Ztg.*, 7, 112.

[2] A. Hilger, 1893: *Vierteljahr. f. öffentl. Gesundpflg*, 25, 559.

ding with those just given for the fourth anhydride, it is perfectly reasonable to suppose that if we had made combustions of a larger number of samples, we might have obtained some results which could be used in support of the idea that these decomposition products of tannins are simple anhydrides of them. Our experience has taught us that the composition of this product is by no means constant with the means used in its preparation, and we saw no good reason for selecting the figures of one combustion from those of a dozen others, presumably as near correct, merely to give support to a theory.

It is somewhat puzzling that Knebel and Hilger have found this body of uniform composition. As a matter of fact Knebel reports no figures in support of his claim, and Hilger reports only two combustions; still it would be supposed that before assigning a formula to a body of this kind a larger number of combustions would have been made, and with different samples. We do not feel justified in accepting the formula $C_{14}H_{13}(OH)_5$ proposed by them for it, nor do we at present possess sufficient data for a formula. We do not even know that it is not a mixture of two or more substances; and although we have not made any separation of them, it seems to us more probable than otherwise that it is a mixture in varying proportions of certain products of decomposition of kolatannin.

The suggestion of Knebel that the tannin of kola is an oxidation product of "kolaroth" according to the equation

$$\underset{\text{Kola red.}}{C_{14}H_{13}(OH)_5} + 4O_2 = \underset{\text{Gallotannic acid.}}{C_{14}H_{10}O_9} + 4H_2O$$

is untenable for the twofold reason that we do not find kolatannin or this decomposition product of it to be so readily oxidizable as stated by him, and furthermore, the kolatannin is not identical with gallotannic acid, the formula of which he has given in the above equation as the formula of kolatannin, nor have we at any time found any gallotannic acid present in kola.

Knebel claims to have prepared a pure product of his kolaroth from the acetyl compound of it; and he further claims that the exact separation of kolaroth may be had by boiling his so-called "kolanin" (caffein kolatannate) with acetyl chloride, whereby an acetyl compound of "kola red" is formed, and is separated

by pouring the mixture of acetyl chloride and caffein kolatannate into water. The acetyl compound of kola red is thus precipitated and the glucose and caffein pass into solution.

Let us examine this statement in the light of the recent knowledge of the chemistry of this interesting drug. First, as conclusively proven by us last year, the body called "kolanin" is not glucoside composed of caffein, glucose and "kola red," but is a combination of caffein with the tannin of kola, $C_{16}H_{20}O_8$, and with anhydrides of that tannin, in varying proportions, depending largely on the method used to separate it from the drug. And it contains no glucose whatever, for we have since learned that the evidence of sugar found in it, and in free kolatannin, last year, and reported in our paper, was due to the accidental presence of glucose from a faulty method of separation. Moreover, acetyl chloride does not effect an exact separation of the caffein from this combination. We prepared an acetyl derivative of caffein kolatannate exactly as done with the other acetyl products and analyzed it, with the following results;

Carbon	58.30
Hydrogen	4.85
Nitrogen	2.54
Oxygen	34.31

The nitrogen content corresponds to 8.79 per cent. of caffein. The caffein kolatannate from which this acetyl compound was made contained 6.53 per cent. nitrogen, corresponding to 22.6 per cent. caffein. Caffein was also found in the liquid from which the acetyl compound was filtered. The separation then is not complete and the acetyl compound is not a pure acetyl compound of caffein kolatannate, nor was it expected to be: for as caffein kolatannate contains kolatannin and some of its anhydrides, both capable of forming acetyl compounds, it naturally follows that any acetyl derivative prepared will not be a simple derivative of one of the bodies, as stated by Knebel, but will be a mixture of acetyl derivatives of the tannin and its anhydrides, probably in the proportions in which they existed prior to the operation. With this understanding of it, it is difficult to see how Knebel could have prepared the pure and simple acetyl compound in the way stated by him.

THE QUESTION OF SUGAR PRODUCTION.

Results were obtained in our work last year which seemed to indicate that kolatannin was a glucosidal body. We wished to investigate this question more fully, and have done so in the following described way :

Kolatannin was dissolved in water and precipitated by lead acetate, and the collected lead salt washed with water for several hours ; the lead salt was then decomposed by dilute sulphuric acid, the lead sulphate filtered out, and the acid filtrate boiled for four hours in a flask fitted with a return condenser. The acid strength was supposed to be about five to six per cent. The insoluble matter was filtered out, the acid removed by addition of barium carbonate, and the liquid again filtered ; the coloring-matter was precipitated by lead acetate and the precipitate removed by filtration, after which the remaining lead acetate was removed by hydrogen sulphide. The filtrate was boiled to remove all the hydrogen sulphide, and made alkaline with a little sodium hydroxide solution. The liquid was still slightly colored. It was tested with Fehling's solution, and gave a red precipitate. A portion of it, made slightly acid with acetic acid, gave negative results when the osazone test was applied, and a control test made at the same time, using a little solution of grape sugar in addition, gave the well-known yellow precipitate. This showed that there was no interference with the phenylhydrazin test. Basic lead acetate solution was added, which precipitated the traces of coloring-matter in the solution, and, after removing the excess of lead, the solution was tested with Fehling's solution again, with negative results this time.

This experiment was repeated in a slightly different way, as follows : The well-washed lead salt of about one gram kolatannin was heated for two hours at 110° C. in a sealed tube with thirty cc. hydrochloric acid, the tube then opened and the contents filtered to remove lead chloride and the amorphous red decomposition product of kolatannin. The filtrate was red-colored and was shaken out several times with ether. The ethereal solutions were set aside for further examination. The filtrate was then boiled to remove the traces of ether, and exactly neutralized with sodium hydroxide, after which basic lead

acetate solution was added until no further precipitation occurred, the precipitate removed by filtration, and the excess of lead salt precipitated as lead sulphate by cautious addition of dilute sulphuric acid. The solution was again filtered, made slightly alkaline with sodium hydroxide solution, and tested with Fehling's solution with negative results.

The importance of using basic lead acetate to precipitate the coloring is shown by the fact that if only normal acetate be used a small amount of coloring-matter is left in solution and gives a precipitate with Fehling's solution, while if basic lead acetate be employed, all the coloring-matter is removed and the copper solution is not affected.

In the ethereal solution, protocatechuic acid in small quantity was found, as stated in another part of this paper.

The experiments described having given negative results, the interpretation is made that kolatannin is not a glucoside and that the evidence we had to the contrary last year was caused by the incomplete removal of the glucose of the plant, due to a faulty method of separation for the tannin, since discarded by us.

It is significant that this result places kolatannin in a category and gives it a character essentially different from the character of caffeotannic acid, as understood by the chemical world from the report of Hlasiwetz.

ACTION OF FUSED ALKALI.

Fifteen grams kolatannin were fused for two hours with potassium hydroxide, the fused mass cooled and acidulated with sulphuric acid, after dissolving it in water. Without filtering the mixture, it was shaken out with ether, the successive portions of this solvent united and evaporated. The residue was distinctly crystalline, the crystals assuming an arborescent form for the most part. This residue was purified several times by recrystallization, after which it was dissolved in water and submitted to tests. Protocatechuic acid and phloroglucin were found to be present.

ACTION OF HEATED GLYCERINE.

One gram kolatannin was heated with three cc. glycerine in a porcelain capsule for twenty minutes at 195°–200°. Fifty cc.

water were then added and the mixture was shaken out with ether. The residue left on evaporating the ethereal extract was found to contain protocatechuic acid, by chemical tests.

KOLATANNIN FROM CAFFEIN KOLATANNATE.

It has thus far not been found practicable to substitute other means of separation of this tannin from its caffein salt for the treatment by lead hydroxide, although the disadvantages of this agent are fully appreciated. It is, however, much to be preferred to lead acetate in this particular case, as lead acetate precipitates some caffein tannate along with the lead tannate, and the resulting tannin is apt to be contaminated accordingly.

PREPARATION OF THE SAMPLE.

The caffein tannate is dried on porous plates and finely powdered and sifted; it is then washed with a mixture of ether and chloroform to remove any traces of adhering fat, and dried again. It is then washed with ice-cold water by agitation, drained and washed again until the washings give no precipitate of silver chloride on being treated with solution of silver nitrate acidulated with nitric acid. The caffein kolatannate is now dissolved in dilute alcohol, making a very concentrated solution; this solution is filtered and poured into ten or twelve times its volume of ice-cold water, whereupon the caffein compound is reprecipitated. It is collected by filtration, drained well and redissolved in dilute alcohol, using gentle heat on the water-bath. Freshly precipitated lead hydroxide triturated to a smooth paste with warm alcohol is now added in excess and the mixture warmed on the water-bath for a few minutes, with stirring. The precipitate is now allowed to subside, and if sufficient lead hydroxide has been added the supernatant liquid will be colorless or faintly straw-colored, while the precipitate of lead kolannate mixed with lead hydroxide forms a compact magma at the bottom of the container. The liquid, which contains kola alkaloids, is poured off and the precipitate is washed repeatedly with dilute alcohol, first by decantation and then by filtration, until several cc. of the filtrate evaporated in a porcelain dish leave no perceptible residue and give no test for caffein. The precipitate is compact and is much more readily washed than that given by lead acetate. It is then

suspended in dilute alcohol and treated with hydrogen sulphide until the tannin is all liberated. The lead sulphide is then filtered out and washed with dilute alcohol until the washings are colorless. The filtrate is concentrated by distillation *in vacuo* to small volume, and eight or ten volumes of water are added. This precipitates a reddish mass, a mixture of certain anhydrides of kolatannin, which is filtered out and set aside. The filtrate containing tannin together with dissolved anhydrides is now shaken with ether once or twice, and the washings thrown away. Then ethyl acetate is added and the tannin separated and purified in the way already described, under the method of preparation of kolatannin.

This "combined" kolatannin, although not quite as light-colored as the "free tannin", shares all its reactions and solubilities.

COMBUSTION OF "COMBINED" KOLATANNIN.

It was analyzed by combustion and the following results obtained, which coincide very closely with those resulting from combustion of the "free" tannin:

	Found.			Calculated for
	I.	II.	III.	$C_{14}H_{20}O_8$.
Carbon	56.53	57.78	56.42	56.45
Hydrogen	5.68	5.79	5.74	5.90
Oxygen	37.79	37.43	37.84	36.65

THE PENTACETYL COMPOUND.

This was prepared in the same manner as the other acetyl derivatives and is identical with the pentacetyl compound of "free" kolatannin, heretofore described.

The analytical results are as follows:

	Found.			Calculated for
By combustion.	I.	II.	III.	$C_{14}H_{15}(C_2H_3O)_5O_8$.
Carbon	56.75	56.44	56.61	56.72
Hydrogen	5.54	5.41	5.31	5.47
Oxygen	37.71	38.15	38.08	37.81
By saponification.				
Acetyl	38.27	38.53	38.68	39.09

THE BROMINE COMPOUNDS.

These were prepared in identically the same way as the corresponding ones of the free tannin. In appearance and properties there seems to be no difference between these and those of

free kolatannin, and for the sake of brevity we shall omit further description and confine ourselves merely to the statement of analytical results in each case.

TRIBROM DERIVATIVE.

	Found.			Calculated for
By combustion.	I.	II.	III.	$C_{16}H_{17}Br_3O_8$.
Carbon	33.09	33.25	....	33.28
Hydrogen	2.84	2.98	....	2.95
Oxygen	22.50	22.13	....	22.19
As silver salt.				
Bromine	41.57	41.64	41.74	41.58

PENTACETYL TRIBROM DERIVATIVE.

(Formed by acetylizing the tribrom derivative.)

	Found.			Calculated for
By combustion.	I.	II.	III.	$C_{16}H_{12}Br_3(C_2H_3O)_5O_8$.
Carbon	39.64	39.82	....	39.65
Hydrogen	3.47	3.61	....	3.44
Oxygen	26.44	25.94	....	26.43
As silver salt				
Bromine	30.45	30.63	30.71	30.48
By saponification.				
Acetyl	26.66	26.97	....	27.32

TRIBROM PENTACETYL DERIVATIVE.

(Formed by brominating the pentacetyl derivative.)

	Found.			Calculated for
By combustion.	I.	II.	III.	$C_{16}H_{12}Br_3(C_2H_3O)_5O_8$.
Carbon	39.88	40.15	....	39.65
Hydrogen	3.26	3.45	...	3.44
Oxygen	26.34	25.74	....	26.43
As silver salt.				
Bromine	30.52	30.66	30.57	30.48
By saponification.				
Acetyl	26.78	27.04	....	27.32

TETRABROM DERIVATIVE.

	Found.		Calculated for
By combustion.	I.	II.	$C_{16}H_{16}Br_4O_8$.
Carbon	29.13	29.35	29.27
Hydrogen	2.29	2.46	2.44
Oxygen	19.86	19.30	19.52
As silver salt.			
Bromine	48.72	48.89	48.77

PENTACETYL TETRABROM DERIVATIVE.

	Found.		Calculated for
By combustion.	I.	II.	$C_{18}H_{11}Br_4(C_2H_3O)_5O_8$.
Carbon	36.12	36.33	36.03
Hydrogen	2.78	2.96	3.01
Oxygen	24.33	23.78	24.02
As silver salt.			
Bromine	36.77	36.93	36.94
By saponification.			
Acetyl	24.64	24.78	24.83

PENTABROM PENTACETYL DERIVATIVE.

(Formed by brominating the pentacetyl derivative.)

	Found.		Calculated for
By combustion.	I.	II.	$C_{18}H_{10}Br_5(C_2H_3O)_5O_8$.
Carbon	33.23	32.98	33.02
Hydrogen	2.62	2.72	2.65
Oxygen	21.71	21.78	22.01
As silver salt.			
Bromine	42.44	42.52	42.32
By saponification.			
Acetyl	22.49	22.26	22.75

HEXABROM DERIVATIVE.

	Found.		Calculated for
By combustion.	I.	II.	$C_{18}H_{14}Br_6O_8$.
Carbon	23.85	23.68	23.59
Hydrogen	1.67	1.83	1.72
Oxygen	14.76	14.72	15.73
As silver salt.			
Bromine	59.62	59.77	58.96

TETRACETYL HEXABROM DERIVATIVE.

	Found.		Calculated for
By combustion.	I.	II.	$C_{18}H_{10}Br_6(C_2H_3O)_4O_8$.
Carbon	29.66	29.92	29.33
Hydrogen	2.46	2.29	2.25
Oxygen	18.73	18.55	19.55
As silver salt.			
Bromine	49.15	49.24	48.87
By saponification.			
Acetyl	16.48	16.73	17.52

KOLATANNIN ANHYDRIDES OBTAINED FROM CAFFEIN KOLATANNATE.

The insoluble, reddish brown precipitate, formed on addition of cold water to the tannin solution after the separation of the tannin from its lead salt, as mentioned in the method of prepa-

ration of "combined" kolatannin, was well washed and dried at 100°. Combustion of this showed a carbon percentage intermediate between that of the second and third anhydrides of kolatannin.

As the solubilities of the anhydrides appear to be about the same, we were unable to effect a separation of them, and for this reason we heated the (supposed) mixture at 135°–140° to constant weight, in order to convert it all if possible into the third anhydride. It was then subjected to another combustion with the following results:

	Found.		Calculated for $(C_{18}H_{17}O_6)_2O$.
	I.	II.	
Carbon	61.27	61.46	61.54
Hydrogen	5.28	5.43	5.13
Oxygen	33.45	33.11	33.33

These results agree very well with the calculated formula and with those obtained by combustion of the third anhydride of so-called "free" tannin of kola.

The identity of the tannin of caffein tannate with the free tannin having been positively settled by the analyses already given, and by one or two evidences yet to be stated, it was deemed unnecessary to make a further comparative study of its bromine derivatives.

ACTION OF DILUTE ACIDS.

The experiments described as having been performed on "free" kolatannin, were repeated on this kind of kolatannin also with parallel results. The same is to be said of the products of fusion with potassium hydroxide and of heating with glycerine.

FURTHER INDICATION OF MOLECULAR CONSTITUTION.

Whether there be a carboxyl in kolatannin or not, remains open to question and to further determinations. Its moderate acid character is not inconsistent with the negative polarity of a polyhydric phenol, reinforced by the oxygen of an anhydride group, yet destitute of entire carboxyl. Our numerous analyses of kolatannin and its several derivatives as described in this paper, are in close agreement with a polyhydric phenol anhydride carrying two methyl groups and one methoxy group, but

standing altogether as a di-addition benzene derivative, two such benzene nuclei pairing together in an ester-like anhydride. On this supposition, a monomethyl dihydrogen trihydroxybenzoic acid has paired ester-wise with a monomethyl methoxydihydrogen trivalent phenol. The di-additive benzene constitution is not improbable among plant constituents of this class, finding parallel in the additive structure of the terpenes. If we admit the hypothesis of the molecular formula above indicated, we must predicate that one of the six hydrogen atoms joined to benzene-carbon is in position somewhat unfavorable to its bromine substitution, and when brominated quite preventing the acetylization of one of the hydroxyls. Finally, there remains no proof as to the distribution of the hydroxyl and methoxyl groups between the two benzene nuclei.

$$\begin{matrix} C_6H_2(CH_3)(OH)_2 & CO \\ & | \\ C_6H_3(CH_3)(OCH_3)(OH)_2 & O \end{matrix} = C_{16}H_{20}O_6.$$

AS TO CHEMICAL BEARINGS UPON PHARMACOLOGY.

Finally, it must be admitted that the stable caffein compound of an acid of such marked chemical individuality as that of kolatannin can by no means be assumed to have the same physiological effects as free caffein. And this subject is one of pharmacological importance for the caffein of all the beverage plants.

We are under many obligations to Mr. James Heggie, B. S., for his very efficient assistance in preparing and analyzing the various derivatives of tannin reported in the preceding pages.

DISCUSSION OF RECENT ASSAY METHODS.

Within the past year two processes for the assay of kola have been offered, which seem to us not to be free from objections. We give below the essential points of these methods without entering into unimportant details.

METHOD OF JEAN.[1]

The author ascribes his plan of procedure in part to MM. Chodat and Chuit. He boils the dried and powdered drug with "milk of lime," dries the whole in an oven and powders it again. This powder is exhausted with chloroform, which

[1] J. Jean, 1896: *Repert. d. Pharm.*, (3), 7, 49.

removes the free alkaloids. The chloroform is evaporated to dryness, the residue dissolved in hot water and filtered. The filtrate is evaporated to dryness and weighed as caffein.

For "kolanin :" the drug, after treatment with chloroform, is extracted with alcohol, the alcohol, evaporated from the percolate and the soft extract remaining is dissolved in boiling water, which solution, after cooling, is filtered. The insoluble matter remaining on the filter is dried in an oven and weighed as "kolanin."

The most serious defects in the above-described method are (1) the boiling with "milk of lime," (2) the prolonged heating necessary to dry the drug after that treatment, (3) weighing the caffein instead of estimating it with Wagner's reagent volumetrically, (4) the means employed to separate so-called "kolanin," (5) estimating caffein kolatannate ("kolanin") by weight as such, instead of by its caffein content.

(1) Boiling with "milk of lime," or other aqueous alkalies, tends to decompose caffein—a fact to well known to require further comment. This treatment also tends to liberate caffein from its tannate, and thus prevents an exact determination of the proportion of "free" to "combined" alkaloid that originally existed in the drug. The starch of kola, amounting to nearly forty per cent.,[1] causes the drug to become of a jelly-like consistence when boiled with aqueous liquids, and when dried the mass is very hard to powder, and even when powdered the condition of it is well calculated to resist the penetrating action of the solvent, and thus cause incomplete extraction.

(2) The gelatinous condition of the drug after boiling with "lime water" renders it hard to dry completely, and as very prolonged heating is necessary, there is danger of loss of caffein by sublimation.

(3) We do not consider that the alkaloids of kola are sufficiently pure when removed in this, the usual manner, to be weighed as such; Gomberg's[2] volumetric method gives more accurate results.

(4) If any caffein kolatannate escapes decomposition during the first part of the essay, which will occasionally happen, it is

[1] Knox and Schlotterbeck, 1895: *Proc. Am. Pharm. Assoc.*, 334.

[2] M. Gomberg, 1896: This Journal, 18, 331.

removed by the extraction with alcohol. But as it is somewhat soluble in water and more soluble in aqueous solutions containing tannin, the directions to collect and weigh the portion left undissolved by water will generally be found superfluous, for the small amount of it present will generally pass into and remain in solution. Dieterich[1] had this experience, and met with nothing but disappointment in trying to estimate caffein kolatannate by this method.

(5) Caffein kolatannate has been shown to be a body of somewhat variable composition, with a caffein content ranging from nineteen to twenty-five per cent. Inasmuch as the value of kola as a stimulant probably depends primarily on its percentage of caffein, and as the weight of its caffein compound indicates only approximately the amount of its combined caffein, there would not seem to be any reason for attaching much importance to the weight of this compound, if a very precise valuation of the drug is desired. We hold that a direct estimation of the caffein of this compound is preferable and at least as expeditious.

METHOD OF CARLES.[2]

Ten grams kola, one gram calcium hydroxide, and twenty grams eighty per cent. alcohol, are mixed together and dried on the water-bath until the weight is reduced to fourteen grams. The mixture is then powdered and transferred to a 100 cc. flask containing thirty-five cc. of a mixture of 100 parts chloroform and twenty parts alcohol, and heated one hour on a water-bath. After filtration the residue is extracted next with twenty cc. of the same solvent, and finally with ten cc. The united extracts are evaporated to dryness, and the residue taken up with ten cc. boiling water containing four or five drops of one per cent. sulphuric acid, then with six cc., and finally with five cc. The solutions are united, filtered, evaporated to constant weight, and weighed as caffein.

For "kolanin," which the author recognizes as caffein kolatannate, another sample of the drug is taken and extracted with water to remove the caffein and other soluble constituents. The drug is next extracted with seventy per cent. alcohol, the extract evaporated to dryness, transferred to a filter, washed

[1] Dieterich, 1896: Chem. Centrbl., 2, 675.
[2] P. Carles, 1896: *J. Pharm. Chim.*, 16, 104.

with cold water, then dried at gentle heat and weighed. If it be desired to estimate the alkaloids of this compound, one gram "kolanin," one gram calcium hydroxide, and three grams chalk with a little seventy per cent. alcohol are mixed together and evaporated on the water-bath to about six grams, and extracted with alcoholic chloroform in the manner already described.

The objections to this method are in part among those already mentioned in the discussion of Jean's method. The use of lime or other alkalies in the assay of a caffein-bearing drug is to be deprecated. The solvent used is not a proper one for the reason that sufficient alcohol is present to extract other constituents in addition to the alkaloids, which are not removed from the caffein during the subsequent treatment of the residue, and which, when weighed with the caffein, lead to erroneous results. Moreover, the manner of applying the menstruum is inconvenient, does not insure complete extraction, and is in no way preferable to the ordinary extraction by the use of Soxhlet's apparatus. The addition of the sulphuric acid is unnecessary and does not add to the purity of the final product, which is dark-colored and very plainly impure. The objection to weighing a final residue as caffein, finds especial application in this method.

As the properties of caffein kolatannate had not been made known very generally at the time of publication of these methods, there is some excuse for the assumption of both of these writers that it is wholly insoluble. Carles has proceeded on this hypothesis in directing the drug to be extracted with cold water to remove the water-soluble constituents before exhausting it with alcohol to remove the caffein compound, but inasmuch as caffein kolatannate is not only somewhat soluble in water, but considerably more soluble in solutions of caffein and of tannin, the extraction of kola by water will remove a considerable amount of it. The same is to be said of the final washing of the caffein compound with water, which is quite inadmissible in quantitative work. Carles seems to have recognized the uncertain value of gravimetric determinations of caffein kolatannate, and is to be commended for offering an alternate method providing for its valuation according to the amount of its alkaloids.

At our request, Mr. James W. Cobb, Ph.C., has assayed a

sample of dried kola by each of the foregoing methods and by the method adopted by us last year,[1] after thoroughly familiarizing himself with them by preliminary work.

	Method of Jean.		Method of Carles.		Knox and Prescott.	
	I.	II.	I.	II.	I.	II.
Caffein by weight.....	1.93	1.79	2.16	2.33	....	....
The same titrated	1.83	1.71	1.87	1.98	1.96	1.88
"Kolanin"	none	none	2.89	3.17	....	....
Combined alkaloids	....	....	....	....	1.62	1.61
Alkaloids of "kolanin"	....	....	0.59	0.69	....	....
Total alkaloids ...	1.83	1.74	2.46	2.67	3.58	3.49

It will be seen that the impurity of caffein separated by Jean's method amounts to 4.84 per cent., and that of the caffein by Carles' method amounts to 14.25 per cent., which was determined by titrating the caffein with Wagner's reagent, after Gomberg's method.

Both give very low results as compared with those obtained by our own. They are tedious and otherwise unsatisfactory, in addition.

Further notes on the assay of kola are in preparation.

All the work reported upon in this paper has been done under the provision of the Stearns Fellowship of the University of Michigan.

ANN ARBOR, MICH., AUGUST, 1897.

NOTE.

A Correction.—In the September number of this Journal the author, together with Mr. F. Thompson, published an article entitled "A Preliminary Thermochemical Study of Iron and Steel," in which they give the results obtained on a number of samples of iron and steel of different composition and which had been subjected to varying heat treatments. On taking up the work again this fall and reviewing critically the results of last year's work, I find that there are at least two serious errors in the results that I wish to explain now in order to prevent anyone else from being misled by the conclusions drawn from the results as stated in the article. The statement that the loss by radia-

[1] Knox and Prescott, 1896: This Journal, 19, 73.

tion was only 0.24° in one hour was a mistake of observation. Careful determinations on this point, made recently, show that it must have been 1.24° instead of the former figure. The error from not taking into account the loss by radiation will amount to from about fifteen to forty calories per gram of metal, the loss depending on the time of solution. Another source of uncertainty, and one for which the author alone is responsible, is the fact that the solution used for dissolving the samples contained about four per cent. of free hydrochloric acid, and consequently the solution of the iron may not in all cases have proceeded according to the equation, $Fe + CuCl_2 = FeCl_2 + Cu$, and this copper redissolved according to the equation $Cu + CuCl_2 = Cu_2Cl_2$. Part of the iron may have dissolved directly in the free hydrochloric acid, and, since the heat absorbed in the decomposition of $2HCl$ is less than that necessary for the reduction of $2CuCl_2$ to Cu_2Cl_2, more heat would be rendered sensible when a good deal of iron was dissolved directly in hydrochloric acid than when solution was effected through cuprous chloride.

The work is being gone over again very carefully with view to eliminating any errors that may exist in the previous work, and I therefore request that judgment be suspended on the previous work until further results may enable us to form a more reliable conclusion than could be drawn from our former data.

E. D. CAMPBELL.

ANN ARBOR, MICH., DECEMBER 14, 1897.

ERRATUM.

On page 942, line 5 (Vol. 19), for "antimony" read "phosphorus.

BOOKS RECEIVED.

A Practical Treatise on Mineral Oils and their By-Products, including a Short History of the Scotch Shale Oil Industry, the Geological and Geographical Description of Scotch Shales, Recovery of Acid and Soda used in Oil Refining and a List of Patents Relating to Apparatus and Processes for Obtaining and Refining Mineral Oils. By Iltyd I. Redwood. London: E. & F. N. Spon, Limited. New York: Spon & Chamberlain. 1897. xiv + 336 pp. Price $6.00

The Early History of Chlorine. Alembic Club Reprints, No. 13. Papers by Carl Wilhelm Scheele (1774), C. L. Berthollet (1785), Guyton de

Morveau (1787), and J. L. Gay-Lussac and L. J. Thenard (1809). Edinburgh: William F. Clay, 1897. 48 pp. Cloth. Price 1s. 6d.

Researches on the Molecular Asymmetry of Natural Organic Products. Alembic Cub Reprints, No. 14. By Louis Pasteur, membre de la Société Chimique de Paris (1860). Edinburgh: William F. Clay. 1897. 46 pp. Cloth. Price 1s. 6d.

Ornithology of North Carolina. A List of the Birds of North Carolina, with Notes of Each Species. Issued by The North Carolina Agricultural Experiment Station, Raleigh, N. C. October 30, 1897. 36 pp.

Current Thought. New Series. Vol. 1, No. 1. January, 1898. C. Elton Blanchard, 802 Ansel Ave., Cleveland, Ohio. 21 pp.

The Mechanics of Soil Moisture. By Lyman J. Briggs, Physicist of the Division of Soils. Bulletin No. 10. U. S. Department of Agriculture, Division of Soils, Washington, D. C. 24 pp.

Second Report on Food Products. Twenty-first Annual Report of the Connecticut Agricultural Experiment Station for 1897. Part I. New Haven: The Tuttle, Morehouse and Taylor Press. 1897. xvii + 64 pp.

Montana Swine Feeding. Bulletin No. 14. Montana Agricultural Experiment Station, Bozeman, Montana. April, 1897. 36 pp. 4 half-tone plates.